

Cambridge Assessment International Education

Cambridge International Advanced Subsidiary and Advanced Level

BIOLOGY 9700/22

Paper 2 AS Level Structured Questions

May/June 2019

MARK SCHEME
Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2019 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of 16 printed pages.



[Turn over

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- · marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

© UCLES 2019 Page 2 of 16

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mark scheme abbreviations

separates marking points

I alternative answers for the same point

R reject

A accept (for answers correctly cued by the question, or guidance for examiners)

I ignore (for answers that include irrelevant information that does not contradict the expected answer)

AW alternative wording (where responses vary more than usual)

ora or reverse argument (for answers which are written as the opposite to the expected answer)

<u>underline</u> actual word given must be used by candidate (grammatical variants accepted)

max indicates the maximum number of marks that can be given

ecf error carried forward

mp marking point (with relevant number)

© UCLES 2019 Page 3 of 16

Question	Answer	Marks
1(a)	stem / not a root, because, transport tissue / vascular bundles / vascular tissue(s) / xylem and phloem / xylem / phloem, around the edges / not in the centre / AW; A stem because, pith / parenchyma tissue, is in the centre	1
1(b)	any two from: (hydrostatic) support / described; R if incorrect context e.g. support because of thick cell walls A packing tissue (cells), turgid / store water; storage; in context of substances other than water e.g. sucrose / starch / waste	2
1(c)(i)	vascular bundle ;	1
1(c)(ii)	0.65 (<u>mm</u>);	1

© UCLES 2019 Page 4 of 16

Question	Answer			
1(d)	mp1-3 must have statements for both xylem and phloem mp4-7 need statements for either xylem or phloem	1		3
	three from:			
	xylem		phloem	
	1 no cytoplasm / hollow / no contents		(peripheral / little / some) cytoplasm; I protoplasm R full of cytoplasm / AW	
	2 no organelles / hollow / no contents		(a few) organelles;A examples of organelles mitochondria or ERI has SER / RER / ribosomesR has nucleus	
	3 lignified R idea of lignin within element		no lignin / (only) cellulose;	
	4 no end wall(s) / no sieve plate(s) A end walls have broken down	or	sieve plate(s) / perforated end wall(s); A (end walls) have sieve pores	
	5 (side walls) contain pits	or	no pits (in side walls);	
	6 no plasmodesmata	or	plasmodesmata (to companion cells);	
	7 thick(er), cell wall / walled	or	thin(ner), cell wall / walled;	

© UCLES 2019 Page 5 of 16

Question		A	nswer		Marks
2(a)	mucous gland cells / mucous gland(s); I mucosa goblet cell(s); list containing trachea and / or bronchus and / or bronchioles I and allow to two max BUT if alveoli stated or incorrect cell types (e.g. ciliated cells) max 1				2
2(b)	any one from: Golgi (body / complex / apparatus); rough endoplasmic reticulum; A rough ER / RER R SER I ER				1
2(c)	two marks to complete table			1	2
		first process	transcription		
		second process	translation		
		third process	glycosylation		
		fourth process	exocytosis		
	allow one mark if transcription first proces allow one mark if transcription and transla				
2(d)	allow one mark if transcription and translation wrong way round and other two correct accept symbol \(\psi\) for water potential chree from: I chloride presence outside cell causes water potential (outside cells) to, decrease / become more negative / become lower or chloride leaving cell causes water potential (in cell) to increase / AW; 2 water potential gradient, created / present (between inside and outside of cell); 3 water leaves cell by osmosis (to be taken up by mucin to form mucus); 4 from higher to lower water potential / AW; R if term gradient included				3

© UCLES 2019 Page 6 of 16

Question	Answer	Marks
2(e)	any three from:	3
	to produce, new / daughter, cells that are genetically identical; to replace, old / damaged / dead, cells; A named cells to repair (damaged / injured / diseased) tissue; A named tissue if mp 2 and 3 not gained, allow 1 mark for replacing damaged tissue	
	 ref.to function; in context AVP; idea of increase in number of cells / growth, to accommodate increase in size of gas exchange system in growing infants / AW 	

Question	Answer	Marks
3(a)	C =neutrophil; A polymorphonuclear leucocyte A basophil A eosinophil D = lymphocyte; B / T is neutral E = monocyte; A macrophage I leucocyte / white blood cell, throughout	3
3(b)(i)	any two from:	2
	lymphocytes / blood cells, produced in / originate from, bone marrow; ref. to stem cell(s); ref. to cell division; e.g. bone marrow has cells that can carry out, mitosis / cell division A (stem cells are) multipotent	

© UCLES 2019 Page 7 of 16

Question	Answer	Marks
3(b)(ii)	any three from:	3
	measles immune response stimulated;	
	detail; e.g. either antigen recognition / presence / binding or lymphocyte / white blood cell, stimulated / triggered plus clonal, expansion / proliferation / cell division / mitosis R incorrect pathogen name R bacteria	
	CLL result of, uncontrolled mitosis; A uncontrolled, division / cell cycle I rapid mitosis	
	detail; owing to, mutation / named mutation or cell cycle checkpoints not working or cells don't respond to signals to control division	
3(c)(i)	any two from:	2
	 (passive / simple) diffusion (of oxygen) across the phospholipid bilayer; A movement from a higher to a lower concentration for diffusion A between phospholipids / between fatty acid chains / across hydrophobic core / across bilayer / AW, for phospholipid bilayer 	
	small size; non-polar / uncharged;	

© UCLES 2019 Page 8 of 16

Question Answer Answer Marks 3(c)(ii) any three from: less oxygen in inhaled air / AW; A less oxygen taken up from alveoli low(er) saturation of haemoglobin (with oxygen); A Hb for haemoglobin A haemoglobin carries less oxygen must be the context of oxygen R saturation, with air / carbon dioxide haemoglobin has lower affinity for oxygen; (so) more haemoglobin required (so more red blood cells) / (more red blood cells) so more haemoglobin; AW compensation / compensates; A so same, volume / quantity / amount, of oxygen reaches body cells (compared to lower altitude)

© UCLES 2019 Page 9 of 16

Question	Answer	Marks
3(d)	any four from:	4
	1 (alleles have) different nucleotide sequences ; A different base sequences	
	2 base substitution / substitution of a base ; A T replaced by A	
	(alleles have) different mRNA codons ; A altered mRNA codon R altered genetic code	
	4 (results in) one amino acid change or change from, glutamic acid / glutamate / glu, to, valine / val or changed primary structure; A changed sequence of amino acids R if implies whole sequence changes	
	5 AVP; e.g. Hb ^A = CTC and Hb ^S = CAC (on template strand) A Hb ^A = GAG and Hb ^S = GTG (on non-template strand) or Hb ^A = CTT and Hb ^S = CAT (on template strand) A Hb ^A = GAA and Hb ^S = GTA (on non-template strand) this is not also mp2 ref. to amino acid substitution at position 6 (this is not also mp4) A 6th, codon / triplet, changed	
	6 changed, (β-globin / Hb) tertiary structure ; A polypeptide / (Hb molecule) quaternary structure R β-globin / polypeptide	
	7 ref. to loss of globular structure / hydrophobic amino acid faces outside (instead of towards centre); A described e.g. (val) amino acid with hydrophobic R-group instead of (glu) amino acid with hydrophilic R-group accept hydrophilic / polar, amino acid accept hydrophobic / non polar, amino acid	

© UCLES 2019 Page 10 of 16

May/June 2019

Question		Answer	Marks
3(d)	8	sticky (haemoglobin) molecule formed / forms fibres with other haemoglobin molecules;	
	9	oxygen taken up less easily / less oxygen carried by haemoglobin / AW ; \pmb{A} $\beta\text{-globin}$ takes up oxygen less easily / AW	
	10	AVP ; (sickle cell) haemoglobin / β -globin, less soluble / AW $_{\mbox{\scriptsize I}}$ insoluble	

© UCLES 2019 Page 11 of 16

Question	Answer	Marks
4(a)	max 3 if any membrane-bound organelles drawn inside cell e.g. nucleus, Golgi body, mitochondrion, lysosome, ER, a large vacuole BUT I vesicle / small vacuoles	4
	I detail of cell wall / cell surface membrane around flagellar area	
	four from: cell wall ; must add another line to diagram R cellulose cell wall	
	cell (surface) membrane ; A plasma membrane label line to the inside of cell wall I if outer line labelled as cell membrane	
	DNA; A bacterial chromosome / nucleoid as label allow one or two circular molecules (circle, loop, ball of string, tangled) R if obviously linear R if label includes, histones / histone proteins / chromatin R if nuclear, envelope / membrane, shown R if nucleolus shown or labelled	
	ribosomes; R 80S ribosomes shown as, dots/small spheres cytoplasm/cytosol;	
	plasmid ; smaller than DNA, circular	
	AVP; e.g. 70S / smaller / 17–20 nm, for ribosome murein / peptidoglycan (for cell wall – allow even if cell wall label not added) pilus /pili drawn as external hair-like structure(s) basal granule at base of flagellum capsule drawn to outside (some do have a capsule)	
4(b)(i)	0.8 % ; R 0.80 %	1

© UCLES 2019 Page 12 of 16

9700/22

Cambridge International AS/A Level – Mark Scheme **PUBLISHED**

Question	Answer	Marks
4(b)(ii)	any two from:	2
	delay in / no, diagnosis; delay in / no, treatment / therapy; A feature of, oral rehydration, treatment /solution A ORT / ORS e.g. lack of supply of ORT unwillingness by health workers to give ORT greater belief in herbal remedies thinking drug treatment is sufficient lack of clean water to make up ORT solution no rehydration programmes I improper treatment / treatment not effective ref. to no / lack of, antibiotic therapy; e.g. lack of supply of / less efficacious (AW) antibiotics used / less supervision in taking full dose ref. to antibiotic resistant strains; idea that already have a weak immune system / malnourished; A may have HIV/AIDS (hence weak immune system)	
4(c)(i)	any one from:	1
	damaged sanitation system / poor sanitation following earthquake ; I poor sanitation unqualified	
	damage to, sewage treatment plants / water purification plants ; AW	
	(contaminated) sewage contaminates drinking water ; I pollution A water becomes contaminated	
	lack of purified drinking water; A lack of, clean / treated / safe / bottled, water	
	AVP ; no, proper / safe, disposal of sewage	

© UCLES 2019 Page 13 of 16

Question	Answer	Marks
4(c)(ii)	any three from:	3
	 2011, peak in / highest / AW, number of, cases / countries / countries and cases; A decrease, increase in 2011, decrease for countries A increase to 2011 and decrease for cases 	
	 (2010 outbreak in) ref. to Haiti and epidemic (so high number of 2011 cases); ref. to spread of disease to countries neighbouring Haiti (hence increase in countries in 2011); overall / generally / AW, decrease in number of countries with cases of cholera or 2008 54–56 countries and 2015 41–43 countries, with ref. to decrease; decrease, peak at 2011 / increase in 2011, decrease is mp1 only 	
	suggestion for trend in decrease in number of countries; e.g. improved infrastructure for sewage / water treatment improved health education to prevent spread providing piped (treated) water	
	 2008–2010 number of cases increased and number of countries decreased; AVP; e.g. lowest / AW, number of cases in 2013 R if also state that 2013 is lowest for countries similar number of countries in 2008 and 2011 but, approx 3×/ much higher / stated values, number of cases in 2011 	

© UCLES 2019 Page 14 of 16

Question		Ans	wer		Marks
5(a)	glycine in monomer column only;				5
	monomer	polymer	monosaccharide	polysaccharide	
	$\begin{array}{c c} \text{(nucleotide)} & \text{m} \\ \alpha\text{-glucose} & \text{;} & \text{g} \end{array}$	cellulose messenger RNA glycogen x-globin;	$\left.\begin{array}{c} \beta\text{-glucose} \\ \alpha\text{-glucose} \end{array}\right\} \; ;$	cellulose ; glycogen ;	
	1	glycine	I glycine	I glycine	
	I incorrect spellings if α-globin appears once and is in monome otherwise correct, allow one mark (ecf)	er column instead of p	polymer column, and the	e monomer and polymer columns a	nre
5(b)	 any three from: allow points from an annotated diagram phosphate head = hydrophilic / polar; fatty acid / hydrocarbon, tail / chains = if mps 1 and 2 not gained, allow one mark 	hydrophobic / non-po		bic / non-polar, tails	3
	3 (so) heads face, watery environment /	tissue fluid / cytoplas	m / cytosol / aqueous er	nvironment;	
	4 (fatty acid) tails, form hydrophobic core	re / form area away fro	om water / face each oth	ner/AW;	
	5 ref. tails and hydrophobic interactions	;			

© UCLES 2019 Page 15 of 16

Question	Answer	Marks
6(a)(i)	any two from:	2
	correct ref. to time; correct ref. to volume of oxygen; A gas for oxygen A volume of water displaced by oxygen A measuring height of water level (decreasing) A cm ³ instead of volume AVP; subtract volume of gas displaced by H ₂ O ₂ addition	
6(a)(ii)	any two from:	2
	greater surface area (over which catalase released);	
	greater number of cells, damaged / cut open, to release catalase; higher concentration of, catalase / enzyme; A more, catalase / enzyme A more active sites available	
	must be linked to idea of more enzyme (i.e. must get mp 2 or 3) so, higher rate of reaction / more enzyme-substrate complexes / more oxygen released; AW	
6(b)	increase to optimum and steeper decrease ; A if curve does not touch one or both axes	1

© UCLES 2019 Page 16 of 16