

## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education

Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		



**CHEMISTRY** 9701/23

Paper 2 Structured Questions AS Core

May/June 2010

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

## **READ THESE INSTRUCTIONS FIRST**

Write your name, Centre number and candidate number on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs, or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE ON ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

The number of marks is given in brackets [ ] at the end of each question or part question.

At the end of the examination, fasten all your work securely together.

For Examiner's Use			
1			
2			
3			
4			
5			
Total			

This document consists of 11 printed pages and 1 blank page.

DC (LEO) 21255/1 © UCLES 2010



[Turn over

## Answer all the questions in the spaces provided.

For
Examiner's
Llco

Hydrazine, N<sub>2</sub>H<sub>4</sub>, can be used as a rocket fuel and is stored as a liquid. It reacts exothermically with oxygen to give only gaseous products.

The enthalpy change of a reaction such as that between hydrazine and oxygen may be calculated by using standard enthalpy changes of formation.

(a) Define the term standard enthalpy change of formation,  $\Delta H_f^{\circ}$ .


(b) Hydrazine reacts with oxygen according to the following equation.

$${\rm N_2H_4(I)} \ + \ {\rm O_2(g)} \ \longrightarrow \ {\rm N_2(g)} \ + \ 2{\rm H_2O(g)}$$

(i) Use the data in the table to calculate the standard enthalpy change of this reaction.

compound	ΔH <sub>f</sub> <sup>⊕</sup> /kJ mol <sup>-1</sup>
N <sub>2</sub> H <sub>4</sub> (I)	50.6
H <sub>2</sub> O(g)	-241.8

$\Delta H^{\circ} =$	 kJ mol <sup>−1</sup>
<b></b>	 11011101

(ii) Although the above reaction is highly exothermic, hydrazine does not burn spontaneously in oxygen.

Suggest a reason for this.

© UCLES 2010 9701/23/M/J/10

	(iii)	Suggest why using hydrazine as a rocket fuel could be regarded as being 'environmentally friendly'.	For Examiner's Use
		[4]	
(c)	The	e bonding in hydrazine is similar to that in ammonia.	
	(i)	Showing outer-shell electrons only, draw a 'dot-and-cross' diagram of an ammonia molecule.	
	(ii)	Draw a diagram to show the three-dimensional shape of an ammonia molecule.	
	(iii)	Draw a diagram to show the shape of a hydrazine molecule. Show clearly which atom is joined to which and show clearly the value of <b>one</b> bond angle.	
		[4]	
(d)	Dec	duce the oxidation state of nitrogen in hydrazine.	
		[1]	
		[Total: 12]	

For Examiner's Use

The alkali metals are a series of six elements in Group I of the Periodic Table. The first ionisation energy of these elements shows a marked trend as the Group is descended.
(a) Define the term first ionisation energy.
[2]
(b) (i) State and explain the trend in first ionisation energy as Group I is descended.
(ii) Suggest how this trend helps to explain the increase in the reactivity of the elements as the Group is descended.
[3]
(c) In a redox reaction, 0.83g of lithium reacted with water to form 0.50 dm <sup>3</sup> of aqueous lithium hydroxide.
$2\text{Li(s)} + 2\text{H}_2\text{O(I)} \rightarrow 2\text{LiOH(aq)} + \text{H}_2(g)$
(i) Calculate the amount, in moles, of lithium that reacted.

© UCLES 2010 9701/23/M/J/10

2

	(ii)	Calculate the volume of hydrogen produced at room temperature and pressure.	For Examiner's Use
	(iii)	Calculate the concentration, in mol dm <sup>-3</sup> , of the LiOH(aq) formed.	
		[5]	
(d)		en heated in chlorine, all of the alkali metals react to form the corresponding bride.	
		cribe what you see when sodium is heated in chlorine and write a balanced equation he reaction.	
	des	cription	
	equ	ation	
		[2]	
		[Total: 12]	

[Turn over © UCLES 2010 9701/23/M/J/10

This o	que	stic	n re	fers	to the	e ele	ment	ts sh	own	in th	e sed	ction	of th	e Pei	riodic	: Tab	le be	low.		For
								Н										He		Examiner's Use
Li	В	е											В	С	N	0	F	Ne		
Na	Μ	g											Αl	Si	Р	S	Cl	Ar		
K	С	а	Sc	Ti	٧	Cr	Mn	Fe	Со	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr		
							ts, id ool of				n cas	se <b>o</b> i	n <b>e</b> e	leme	nt th	at h	as th	ne pro	operty	
(	i)	an	ele	ment	that	sink	s in c	old v	wate	and	read	cts re	adily	with	it					
(i	i)	an	مام	ment	that	form	ns an	oxid	le tha	ıt iç a	a red	ucina	n ane	nt						
(1	''	an	CIC	illelli	ınaı	10111	is an	OXIO		11 13 6	ı icu	uciri	y age	,111						
				•••																
(ii	i)	the	e ele	men	t tha	t has	the	large	est fir	st io	nisati	on e	nerg	y						
(iv	Λ	the	me	ıtal ir	n Per	iod :	3 (Na	to A	r) the	nt ha	s the	ema	llest	catio	'n					
(1)	,	tiic	, 1110	idi ii	11 01	100 0	) (I <b>V</b> a	10 / 1	1) 1110	tt Ha	5 1110	Silic	illoot	cano	'11					
				•••																
(1	/)						as a ar stri	_		lecul	ar st	ructu	ıre <b>a</b> ı	<b>nd</b> fo	rms	an o	xide	whic	h also	
(v	i)	the	e ele	men	t in F	Perio	1) & b	Na to	Ar)	with	the g	great	est e	lectri	cal c	ondu	ctivi	ty		
																			[6]	
							Perio weer					ident	ify <b>tv</b>	<b>ио</b> е	leme	ents v	whos	se hyd	drides	
			an	d															[1]	

© UCLES 2010 9701/23/M/J/10

3

For Examiner's Use

. ,	iden	the elements in Period 3 (Na to Ar) in the section of the Periodic Table opposite to tify the oxide(s) referred to below. <b>ach</b> case, give the formula of the oxide(s).
	(i)	an oxide which has no reaction with water
. (	(ii)	two acidic oxides formed by the same element
		and
(1	iii)	an oxide which dissolves readily in water to give a strongly alkaline solution
(	iv)	an oxide which is amphoteric
		[5]
		[Total: 12]

8 4 Organic reactions involve substances which may be atoms, molecules, ions or free radicals. We also apply the terms electrophilic, nucleophilic, addition, elimination and substitution to organic reactions. Consider the following reactions.  $CH_4 + Cl_2 \rightarrow CH_3Cl + HCl$ reaction 1  $CH_3CH_2OH \rightarrow CH_2=CH_2 + H_2O$ reaction 2  $CH_3I + OH^- \rightarrow CH_3OH + I^$ reaction 3  $CH_3COCH_3 + HCN \rightarrow CH_3C(OH)(CN)CH_3$ reaction 4 (a) Using the terms mentioned above, state as clearly as you can the nature of each of the following reactions. reaction 1 ..... [2] reaction 2 ..... (b) By considering the four reactions above, suggest a formula for each of the following substances. In **each** case, state which reaction you are considering. (i) one substance that is an addition product reaction..... addition product (ii) one substance that is a leaving group reaction..... leaving group

[3]

For Examiner's Use

© UCLES 2010 9701/23/M/J/10

reaction.....

(iii) one substance that behaves as an electrophile

electrophile

(c)	What is meant by the term <i>nucleophile</i> ?	For Examiner's
		Use
	[1]	
(d)	Reactions 3 and 4 involve nucleophiles.	
	For <b>each</b> reaction, give the formula of the nucleophile.	
	reaction 3	
	reaction 4	
	[2]	
(e)	One characteristic reaction of ethene is its ability to decolourise bromine.	
	$CH_2=CH_2 + Br_2 \rightarrow BrCH_2CH_2Br$	
	In this reaction, ethene behaves as a nucleophile.	
	Suggest an explanation for how ethene can behave in this way.	
	[1]	
	[Total: 9]	

5	Lactic acid, 2-hydroxypropanoic acid, CH <sub>3</sub> CH(OH)CO <sub>2</sub> H, occurs naturally in sour milk and in
	our muscles when we take hard exercise.
	Lactic acid is chiral and shows stereoisomerism.

For Examiner's Use

(a) Draw fully displayed structures of the two optical isomers of lactic acid. Indicate with an asterisk (\*) the chiral carbon atom in the lactic acid molecule.

[3]

**(b)** Lactic acid may be synthesised from ethanol by the following route.

$$\mathsf{CH_3CH_2OH} \xrightarrow{\mathsf{step 1}} \mathsf{CH_3CHO} \xrightarrow{\mathsf{step 2}} \mathsf{CH_3CH(OH)CN} \xrightarrow{\mathsf{step 3}} \mathsf{CH_3CH(OH)CO_2H}$$

Give the reagent(s) and essential condition(s) for **each** step.

	reagent(s)	condition(s)
step 1		
step 2		
step 3		

[6]

During exercise, lactic acid is produced in our muscles from pyruvic acid, $\mathrm{CH_3COCO_2H}$ . This reaction occurs in the presence of the enzyme lactic acid dehydrogenase.					
(c) (i)	What type of chemical compound is the enzyme lactic acid dehydrogenase?				
(ii)	How would you detect a small quantity of pyruvic acid in a sample of lactic acid?				
	State the reagent(s) you would use and what would be seen in your test.				
	reagent(s)				
	observation				
(iii)	How would you detect a small quantity of lactic acid in a sample of pyruvic acid?				
	State the reagent(s) you would use and what would be seen in your test.				
	reagent(s)				
	observation				
(iv)	What chemical reagent would be used to convert pyruvic acid into lactic acid?				
	$CH_3COCO_2H \rightarrow CH_3CH(OH)CO_2H$				

[Total: 15]

[6]

For Examiner's Use

© UCLES 2010 9701/23/M/J/10

## **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2010 9701/23/M/J/10