MARK SCHEME for the October/November 2010 question paper

for the guidance of teachers

9701 CHEMISTRY

9701/35

Paper 3 (Advanced Practical Skills), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A / AS LEVEL – October/November 2010	9701	35

Questior	n Sections	Indicative material	Mark	
1 (a)	PDO Layout	 Volume given for Rough titre. and accurate titre details tabulated. 	1	
	MMO Collection	 Follows instructions – initial and final burette readings recorded for Rough titre and initial and final burette readings and volume of FA 2 added recorded for each accurate titre and headings should match readings. Do not award this mark if: 50(.00) is used as an initial burette reading; more than one final burette reading is 50.(00); any burette reading is greater than 50.(00) 	1	
	MMO Decisions	 III Has two uncorrected, accurate titres within 0.1 cm³ Do not consider the Rough even if ticked. Do not award this mark if having performed two titres within 0.1 cm³ a further titration is performed which is more than 0.10 cm³ from the closer of the initial two titres, unless a fourth titration, within 0.1 cm³ of the third titration has also been carried out. 	1	
	PDO Recording	IV All accurate burette readings (initial and final) recorded to nearest 0.05 cm ³ Assess this mark on burette readings only	1	
	MMO Quality	 V, VI and VII Round any burette readings to the nearest 0.05 cm³. Check and correct subtractions in the titre table. Select the "best" titre using the hierarchy: two identical; titres within 0.05 cm³; titres within 0.1 cm³; etc. Award <u>V, VI and VII</u> for a difference from Supervisor within 0.20 cm³ 	3	
		Award <u>V</u> and VI only for a difference of 0.20+ cm ³ – 0.30 cm ³		
		Award <u>V only</u> for a difference of 0.30+ - 0.50 cm ³ If the "best" titres are ≥ 0.50 cm ³ apart cancel one of the Q marks.		[7]

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A / AS LEVEL – October/November 2010	9701	35

(b)	ACE Interpretation	Calculates the mean, correct to 2 decimal places from any accurate titres within 0.20 cm ³ . The third decimal place may be rounded to the nearest 0.05 cm^3 . A mean of exactly .x25 or .x75 is allowed but the candidate may round up or down to the nearest 0.05 cm ³ . If ALL burette readings are given to 1 decimal place then the mean can be given to 1 decimal place if numerically correct without rounding. Mean of 24.3 and 24.4 = 24.35 (\checkmark) Mean of 24.3 and 24.4 = 24.4 (\star) Titres to be used in calculating the mean must be	1	
		clearly shown – in an expression or ticked in the titration table.		[1]
(c)	ACE Interpretation	I Correctly evaluates $\frac{10.00}{40} = 0.25(0)$	1	
		II Uses answer (i) $\times \frac{\text{mean titre}}{1000}$ in step (ii)	1	
		and 1000		
		answer (ii) x ¹⁰⁰⁰ /10 in step (iii)		
		If an answer, with no working, is given in any section allow if correct.		[2]
	Total		[Tot	al: 10]

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A / AS LEVEL – October/November 2010	9701	35

2 (a)	PDO Recording	I Has correct headings (minimum three) and units in the weighing table in (2)(a) and correct units in the titration table in (2)(b)	1	
		Acceptable units are /g, (g), mass in grams, mass in g; similarly /cm³,		
		 All three balance reading are read with constant precision (same no of decimal places) and to at least 1 decimal place 	1	[2]
Calculate 8 Mass of FA	× (3.00 – mass 3 used = (mass	he titre for 3.00 g of FA 3 added to the acid. of FA 3 used) and subtract from the titre obtained. tube + FA 3) – (mass tube + residue) hass of empty tube then use (mass tube + FA 3) – (mas	s tube).	
(b)	MMO Quality	Award <u>I and II</u> if the difference between candidate and Supervisor scaled titres is within 0.40 cm^3	1	
		Award <u>I only</u> if the difference is between 0.50+ cm^3 and 0.80 cm^3	1	[2]
(c)	There is no m	ark available for this section.		
(d)	ACE Interpretation	I Uses $\frac{\text{mean titre}}{1000} \times 0.280$ in step (i)	1	
		and uses answer (i) $\times \frac{250}{25}$ in step (ii)		
		II Correctly evaluates $\frac{0.5 \times 250}{1000} = 0.125$ in step	1	
	PDO Display	III Uses answer (iv) × 0.5 × 100 in step (v)	1	
		 IV Working shown in a minimum of three sections Working should be a step in the right direction: step (i) 0.28 × titre volume (in cm³/dm³) step (ii) Use of 25 & 250 or 10 step (iii) 0.5 and 250 step (iv) the correct two numbers step (v) would need to include 2 (0.5) and 100 step (vi) must be correct 	1	
		 V 3 to 5 significant figures in final answers to all sections attempted – minimum of three final answers required 	1	[5]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A / AS LEVEL – October/November 2010	9701	35

	1		1	i
(e)	ACE Conclusions	Explains one of the following: If 5.5 g of CaCO ₃ had been used the titre would be too small/not enough HC <i>l</i> remains for the <u>titration</u> (not 'all the acid has reacted') or Difficult/takes too long to dissolve 5.5 g of solid/it will not all dissolve <u>in 150 cm³</u> (of acid) or Excessive/too fast effervescence/fizzing/rate of gas evolved or	1	
		Acid spray		[1]
(f)	ACE Interpretation	 (i) If balance displays to 1 decimal place: error in balance reading is ±0.05 g or ±0.1(0) g error in mass of FA 3 is ±0.1 g or ±0.2 g If balance displays to 2 decimal places: error in balance reading is ±0.005 g or ±0.01 g error in mass of FA 3 is ±0.01 g or ±0.02 g If balance displays to 3 decimal places: error in balance reading is ±0.0005 g or 0.001g error in mass of FA 3 is ±0.001 g or ±0.02 g If balance displays to 3 decimal places: error in balance reading is ±0.0005 g or 0.001g error in mass of FA 3 is ±0.001 g or ±0.002 g 	1	
		 (ii) Correctly evaluates to at least 2 significant figures: <u>candidate's error in mass of FA 3</u> × 100 	1	
		mass of FA 3 used		[2]
(g)	ACE Conclusions	 Gives correct equation for the thermal decomposition of calcium carbonate including state symbols 	1	
	ACE Improvements	(ii) Outlines: weigh container weigh container + solid (heating and) weighing again repeated (heating and) weighing to constant mass or weigh container weighing container + solid (heating and) measuring gas volume when no further increase and cooled to room temperature / use of pV = nRT / $\frac{PV}{T}$ = constant	1	[2]
	Total			[14]
	Total			[14]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A / AS LEVEL – October/November 2010	9701	35

		FA7 is Fe ₂ (SO_4) ₃ (aq); FA 8 is $CrCl_3$ (aq); FA 9 is ZnI_2 (aq) [$ZnCl_2 + KI$]		
3	(a)	PDO Layout	 I (Tabulates) observations clearly, showing: observation when each reagent is first added and observation when reagent added to <u>excess</u> (if there is a ppt) 	1	
		MMO Collection	 II, III and IV 1 mark for correct observations in each of the columns or rows representing FA 7, FA 8 and FA 9 or 1 mark for correct observations in the row or column representing a reagent added (initial and excess count as one row/column) 	3	
		ACE Conclusions	Award <u>V only</u> if one ion only is correctly identified	1	
			Award <u>V</u> and <u>VI</u> if all three ions are correctly identified from candidate's observations. Allow ecf*	1	[6]

Minimum for observations marks:

Solution	FA 7	FA 8	FA 9
NaOH	red-brown/brown/rust ppt insoluble (in excess)	grey-green ppt <u>soluble</u> /dissolves (in excess) giving a dark green solution	White/milky white ppt soluble/dissolves (in excess)
NH ₃	red-brown ppt insoluble (in excess) <i>(suitable qualified brown)</i>	grey-green ppt insoluble (in excess)	White/milky white ppt soluble/dissolves (in excess)

Minimum for conclusions marks: (with incomplete but not CON observations)

- **FA** 7 red-brown ppt with either;
- grey-green ppt with either/(dark) green solution with excess NaOH; FA 8
- **FA 9** white ppt soluble in excess NH₃.

* ecfs allowed

- **FA 8** allow Fe^{2+} if green ppt insoluble in excess NaOH (no grey-green ppts) **FA 9** allow Al^{3+} and Pb^{2+} if white ppt insoluble in excess NH₃

- **FA 9** allow Ba^{2+} and NH_4^+ if no ppt with either **FA 9** allow Mg^{2+} if white ppt insoluble in excess of both

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A / AS LEVEL – October/November 2010	9701	35

(b)	MMO Decisions	I	Selects barium chloride or barium nitrate for the test in step (i) Do not allow Ba^{2+} alone $Ba^{2+}(aq)$ or soln containing Ba^{2+} (ions) is acceptable	1	
	MMO Collection	II	Records white/off-white precipitate with only FA 7	1	
	MMO Decisions	111	Selects silver nitrate or lead nitrate in (ii) to add to the solutions (that do not contain sulfate) Do not allow Ag^+ or Pb^{2+} alone Aqueous ions or solutions containing the ion are acceptable as above	1	
	MMO Collection	IV	Appropriate observations FA 8 white ppt with Ag ⁺ /white ppt or no ppt with Pb ²⁺ FA 9 yellow ppt with either <i>Ignore observations with any solution candidate</i> <i>has identified as sulfate</i>	1	
	ACE Conclusions	V	FA 8 is chloride, FA 9 is iodide Credit if the supporting evidence fits the ion identified and the practical performed for FA 8 and FA 9 provided there is no CON observation in (i) <i>Do not credit if Ag</i> ⁺ <i>gives a ppt with FA 7</i> Marks IV and V may be awarded from FA 8 white ppt chloride (IV) FA 9 yellow ppt iodide (V)	1	[5]

Other possibilities:

Two white ppts with aqueous Ba^{2+} then remaining solution tested with aqueous Ag^+/Pb^{2+} This would score marks I, III and may score one of IV or V

Aqueous Ba^{2+} gives positive result with solution other than **FA 7** and tests with aqueous Ag^{+}/Pb^{2+} performed

(This would score marks I and III)

Ignore observation and conclusion with FA 7

Award correct observation and valid conclusion for third ion thus scoring one of IV or V

Aqueous Ba²⁺ gives positive result with all three solutions

Award mark I, and mark III may be awarded for selection of aqueous Ag⁺/Pb²⁺ or statement that no further testing is required **but no other marks can be awarded** in this section.

Page 8	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A / AS LEVEL – October/November 2010	9701	35

	FA 10 is NaNO ₃ (s); FA 11 is NaNO ₂ (s)							
(c) (i)	MMO Collection	I Solid/ FA 10 melts/to a liquid/solution (on heating)	1					
	Concetion	II Observes <u>bubbles</u> of gas in liquid/solution or	1					
		Liquid/solution turns yellow/pale yellow						
	MMO Decisions	III Describes an appropriate test in either (i) or (ii) for any of the following gases: O ₂ , CO ₂ , NH ₃ or SO ₂	1					
		There must be a reference to gas being evolved before this mark can be awarded.						
	MMO Collection	 IV Positive identification of oxygen gas in (i): glowing splint rekindles/relights/glows brighter (gas evolved rekindles a glowing splint would gain marks III and IV) ('glowing splint rekindles' would gain mark III not IV) 	1					
(ii)		V On adding acid to residue to FA 11 , observes brown/yellow-brown gas (<i>not yellow</i> , <i>orange or red-brown</i>) or	1					
		blue solution (allow greenish blue)		[5]				
	Total			[16]				