



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/33

Paper 3 Further Mechanics

May/June 2020

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 ms^{-2} .

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Blank pages are indicated.

- 1 A particle P of mass m is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O on a smooth horizontal plane. The particle P moves in horizontal circles about O . The tension in the string is $4mg$.

Find, in terms of a and g , the time that P takes to make one complete revolution. [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 2 A particle Q of mass m kg falls from rest under gravity. The motion of Q is resisted by a force of magnitude mkv N, where $v \text{ ms}^{-1}$ is the speed of Q at time t s and k is a positive constant.

Find an expression for v in terms of g , k and t . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of horizontal dotted lines for writing.

- 3 A particle Q of mass m is attached to a fixed point O by a light inextensible string of length a . The particle moves in complete vertical circles about O . The points A and B are on the path of Q with AB a diameter of the circle. OA makes an angle of 60° with the downward vertical through O and OB makes an angle of 60° with the upward vertical through O . The speed of Q when it is at A is $2\sqrt{ag}$.

Given that T_A and T_B are the tensions in the string at A and B respectively, find the ratio $T_A : T_B$. [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

.....




A uniform solid circular cone, of vertical height $4r$ and radius $2r$, is attached to a uniform solid cylinder, of height $3r$ and radius kr , where k is a constant less than 2. The base of the cone is joined to one of the circular faces of the cylinder so that the axes of symmetry of the two solids coincide (see diagram). The cone and the cylinder are made of the same material.

- (a)** Show that the distance of the centre of mass of the combined solid from the vertex of the cone is $\frac{(99k^2 + 96)r}{18k^2 + 32}$. [4]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

5



Two uniform smooth spheres A and B of equal radii each have mass m . The two spheres are each moving with speed u on a horizontal surface when they collide. Immediately before the collision A 's direction of motion makes an angle of α° with the line of centres, and B 's direction of motion is perpendicular to that of A (see diagram). The coefficient of restitution between the spheres is e .

Immediately after the collision, B moves in a direction at right angles to the line of centres.

- (a) Show that $\tan \alpha = \frac{1+e}{1-e}$. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....

.....

7



One end of a light spring of natural length a and modulus of elasticity $4mg$ is attached to a fixed point O . The other end of the spring is attached to a particle A of mass km , where k is a constant. Initially the spring lies at rest on a smooth horizontal surface and has length a . A second particle B , of mass m , is moving towards A with speed $\sqrt{\frac{4}{3}ga}$ along the line of the spring from the opposite direction to O (see diagram).

The particles A and B collide and coalesce. At a point C in the subsequent motion, the length of the spring is $\frac{3}{4}a$ and the speed of the combined particle is half of its initial speed.

(a) Find the value of k . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

At the point C the horizontal surface becomes rough, with coefficient of friction μ between the combined particle and the surface. The deceleration of the combined particle at C is $\frac{9}{20}g$.

(b) Find the value of μ . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.