



Cambridge International AS & A Level

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/42

Paper 4 Further Probability & Statistics

October/November 2022

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **12** pages.

- 2 In the colleges in three regions of a particular country, students are given individual targets to achieve. Their performance is measured against their individual target and graded as ‘above target’, ‘on target’ or ‘below target’. For a random sample of students from each of the three regions, the observed frequencies are summarised in the following table.

		Region			
		<i>A</i>	<i>B</i>	<i>C</i>	Total
Performance	Above target	62	41	44	147
	On target	102	94	95	291
	Below target	56	45	61	162
Total		220	180	200	600

Test, at the 10% significance level, whether performance is independent of region. [7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- 3 A scientist is investigating the masses of birds of a certain species in country X and country Y . She takes a random sample of 50 birds of this species from country X and a random sample of 80 birds of this species from country Y . She records their masses in kg, x and y , respectively. Her results are summarised as follows.

$$\Sigma x = 75.5 \quad \Sigma x^2 = 115.2 \quad \Sigma y = 116.8 \quad \Sigma y^2 = 172.6$$

The population mean masses of these birds in countries X and Y are μ_x kg and μ_y kg respectively.

Test, at the 5% significance level, the null hypothesis $\mu_x = \mu_y$ against the alternative hypothesis $\mu_x > \mu_y$. State your conclusion in the context of the question. [8]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A series of 30 horizontal dotted lines for writing.

4 The continuous random variable X has probability density function f given by

$$f(x) = \begin{cases} k & 0 \leq x < 1, \\ kx & 1 \leq x \leq 2, \\ 0 & \text{otherwise,} \end{cases}$$

where k is a constant.

(a) Show that $k = \frac{2}{5}$. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Find the interquartile range of X . [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

5 A 6-sided dice, A , with faces numbered 1, 2, 3, 4, 5, 6 is biased so that the probability of throwing a 6 is $\frac{1}{4}$. The random variable X is the number of 6s obtained when dice A is thrown twice.

(a) Find the probability generating function of X . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

A second dice, B , with faces numbered 1, 2, 3, 4, 5, 6 is unbiased. The random variable Y is the number of 6s obtained when dice B is thrown twice.

The random variable Z is the total number of 6s obtained when both dice are thrown twice.

(b) Find the probability generating function of Z , expressing your answer as a polynomial. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(c) Find $\text{Var}(Z)$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(d) Use the probability generating function of Z to find the most probable value of Z .

[1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(b) Explain whether a paired sample t -test would be appropriate to test the manager's claim if earnings are normally distributed. [1]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

