
PHYSICS

9702/23

Paper 2 AS Level Structured Questions

October/November 2017

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

© IGCSE is a registered trademark.

This document consists of **7** printed pages.

Question	Answer	Marks
1(a)(i)	work (done) / time (taken) or energy (transferred) / time (taken)	B1
1(a)(ii)	Correct substitution of base units of all quantities into any correct equation for power. Examples: $(P = E / t \text{ or } W / t \text{ gives}) \text{ kg m}^2 \text{ s}^{-2} / \text{s} = \text{kg m}^2 \text{ s}^{-3}$ $(P = Fs / t \text{ or } mgh / t \text{ gives}) \text{ kg m s}^{-2} \text{ m} / \text{s} = \text{kg m}^2 \text{ s}^{-3}$ $(P = \frac{1}{2}mv^2 / t \text{ gives}) \text{ kg (m s}^{-1})^2 / \text{s} = \text{kg m}^2 \text{ s}^{-3}$ $(P = Fv \text{ gives}) \text{ kg m s}^{-2} \text{ m s}^{-1} = \text{kg m}^2 \text{ s}^{-3}$ $(P = VI \text{ gives}) \text{ kg m}^2 \text{ s}^{-2} \text{ A}^{-1} \text{ s}^{-1} \text{ A} = \text{kg m}^2 \text{ s}^{-3}$	A1
1(b)(i)	units of A : m^2 and units of T : K	C1
	units of k : $\text{kg m}^2 \text{ s}^{-3} / \text{m}^2 \text{ K}^4$ $= \text{kg s}^{-3} \text{ K}^{-4}$	A1
1(b)(ii)	curve from the origin with increasing gradient	B1

Question	Answer	Marks
2(a)	$\rho = m / V$ or $\rho = m / Ah$	B1
	$p = F / A$ or $p = W / A$	B1
	$p = [\rho Ahg] / A$ or $p = [\rho Vg] / [V / h]$ (so) $p = \rho gh$	A1

Question	Answer	Marks
2(b)(i)	1. weight/gravitational (force) upthrust (force)/buoyancy (force) drag/viscous/frictional (force)/fluid resistance/resistance	B1
	2. weight = upthrust + viscous (force)	B1
2(b)(ii)	<ul style="list-style-type: none"> • decrease in (gravitational) potential energy (of sphere) due to decrease in height (since $E_p = mgh$) • increase in thermal energy due to work done against viscous force/drag • loss/change of (total) E_p equal to gain/change in thermal energy Any 2 points.	B2
2(c)(i)	atmospheric pressure = $9.1(0) \times 10^4$ Pa	A1
2(c)(ii)	$(\Delta)p = \rho g(\Delta)h$ $(9.15 - 9.10) \times 10^4 = \rho \times 9.81 \times (0.17 - 0.10)$	C1
	$\rho = 730$ (728) kg m^{-3}	A1

Question	Answer	Marks
3(a)	<u>sum/total</u> momentum (of system of bodies) is constant or <u>sum/total</u> momentum before = <u>sum/total</u> momentum after	M1
	for an isolated system/no (resultant) <u>external</u> force	A1
3(b)(i)	$p = mv$	C1
	$(4.0 \times 6.0 \times \sin \theta) - (12 \times 3.5 \times \sin 30^\circ) = 0$ or $(m_A v_A \times \sin \theta) - (m_B v_B \times \sin 30^\circ) = 0$	M1
	$\theta = 61^\circ$	A1

Question	Answer	Marks
3(b)(ii)	shows the horizontal <u>momentum</u> component of ball A or of ball B as $(4.0 \times 6.0 \times \cos \theta)$ or $(12 \times 3.5 \times \cos 30^\circ)$	C1
	$(4.0 \times 6.0 \times \cos 61^\circ) + (12 \times 3.5 \times \cos 30^\circ) = 4.0v$ so $v = 12 \text{ (ms}^{-1}\text{)}$	A1
3(b)(iii)	initial $E_K (= \frac{1}{2} \times 4.0 \times 12^2) = 290 \text{ (288) (J)}$	M1
	final $E_K (= \frac{1}{2} \times 4.0 \times 6.0^2 + \frac{1}{2} \times 12 \times 3.5^2) = 150 \text{ (145.5) (J)}$	M1
	(initial $E_K >$ final E_K) so inelastic [both M1 marks required to award this mark]	A1

Question	Answer	Marks
4(a)	displacement of particles/vibration(s)/oscillation(s) is parallel to/along the direction of energy/propagation	B1
4(b)	period = $1 / 800 (= 1.25 \times 10^{-3} \text{ s})$	C1
	time-base setting = $1.25 \times 10^{-3} / 2.5$	C1
	$= 5.0 \times 10^{-4} \text{ s cm}^{-1}$	A1
4(c)(i)	$I \propto A^2$	C1
	$(I_X / I_Y =) [r_Y / r_X]^2 = [A_X / A_Y]^2$	C1
	ratio $A_Y / A_X = 120 / 30$ $= 4.0$	A1

Question	Answer	Marks
4(c)(ii)	1. $v = f\lambda$	C1
	minimum $\lambda = 330 / (800 + 16) = 0.40 \text{ m}$	A1
	2. $f_o / f_s = v / (v - v_s)$ $816 / 800 = 330 / (330 - v_s)$	C1
	$v_s = 6.5 \text{ m s}^{-1}$	A1

Question	Answer	Marks
5(a)	force <u>per</u> unit positive charge	B1
5(b)(i)	$s = \frac{1}{2}at^2$	C1
	$a = (2 \times 0.045) / (1.5 \times 10^{-7})^2 = 4(.0) \times 10^{12} \text{ m s}^{-2}$	A1
5(b)(ii)	$F = 1.67 \times 10^{-27} \times 4.0 \times 10^{12} = 6.7 (6.68) \times 10^{-15} \text{ N}$	A1
5(b)(iii)	1. $E = F / Q$	C1
	$= 6.68 \times 10^{-15} / 1.6 \times 10^{-19}$ $= 4.2 (4.18) \times 10^4 \text{ NC}^{-1}$	A1
	2. $E = V / d$	C1
	$V = 4.18 \times 10^4 \times 0.045$ $= 1.9 \times 10^3 \text{ V}$	A1

Question	Answer	Marks
5(c)	$a = Eq / m$ or $F = ma$ and $F = Eq$	C1
	ratio = $\frac{(2 \times 1.6 \times 10^{-19}) \times (1.67 \times 10^{-27})}{(1.6 \times 10^{-19}) \times (4 \times 1.66 \times 10^{-27})}$ or $\frac{2 \times 1}{1 \times 4}$ = 0.50	A1

Question	Answer	Marks
6(a)(i)	$P = VI$	C1
	$I = 30 / 120$ = 0.25 A	A1
6(a)(ii)	$Q = 0.25 \times 3.0 \times 3600 (= 2700)$	C1
	number = $(0.25 \times 3.0 \times 3600) / 1.60 \times 10^{-19}$ = 1.7×10^{22}	A1
6(b)	$R = V/I$ or $R = P/I^2$ or $R = V^2/P$	C1
	= $120 / 0.25$ or = $30 / 0.25^2$ or = $120^2 / 30$ = 480 Ω	A1

Question	Answer	Marks
6(c)	$R = \rho l / A$	C1
	$A = (6.1 \times 10^{-7} \times 580 \times 10^{-3}) / 480 (= 7.37 \times 10^{-10})$	C1
	$d = [(4 \times 7.37 \times 10^{-10}) / \pi]^{1/2}$ $= 3.1 \times 10^{-5} \text{ m}$	A1
6(d)	temperature decreases and so resistance decreases	B1

Question	Answer	Marks
7(a)	nucleons = 23	B1
	neutrons = 11	B1
7(b)	similarity: same (rest) mass or equal (magnitude of) charge	B1
	difference: opposite (sign of) charge or one is matter and one is antimatter or one is an electron and one is an antielectron	B1