

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME		
CENTRE NUMBER		CANDIDATE NUMBER
CHEMISTRY		0620/62
Paper 6 Alterna	tive to Practical	May/June 2012
		1 hour
Candidates ans	wer on the Question Paper.	
No Additional M	aterials are required.	

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.


At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
4		
5		
6		
7		
Total		

This document consists of 12 printed pages.



1 A student heated red crystals of hydrated cobalt(II) chloride, $CoCl_2.6H_2O$, to obtain a sample of water. He used two different sets of apparatus, **A** and **B**.



(a) Complete the box to identify the piece of apparatus labelled.

[1]

For

Examiner's

Use

(b) The steam could have been condensed more efficiently using a condenser. Draw a labelled diagram of a condenser.

https://xtremepape.rs/

(c)	The colour of the solid remaining in the boiling tube after heating was blue. Predict the effect of adding water to this solid. Explain your answer.	For Examiner's Use
	effect	
	explanation[2]	
(d)	Suggest why the boiling tube cracked using set of apparatus B but not set A .	
	[2]	
	[Total: 7]	


3

For Examiner's Use

2 An experiment was carried out to measure the speed (rate) of reaction between magnesium carbonate and excess dilute nitric acid. 50 cm³ of dilute nitric acid was poured into a conical flask and placed on a balance. 1.0 g of powdered magnesium carbonate was added to the flask. The mass of the flask and contents decreased as a gas was given off. The loss in mass was recorded every half minute for six

The results of the experiment are shown plotted on the grid below.

minutes.



3 Electricity was passed through a solution of concentrated hydrochloric acid as shown below.

For Examiner's Use

Bubl	bles	were observed at both electrodes.
(a)	Giv	e one other expected observation.
(b)	Lab	el the electrodes. [1]
(c)	(i)	Name the gas given off at the cathode (negative electrode).
		[1]
((ii)	Give a test for this gas.
		test
		result[2]

For A student investigated a reaction between a solid and a liquid. The reaction produced a gas. 4 Examiner's She wanted to know if any of the substances **W**, **X** and **Y** were catalysts for the reaction. Use Firstly she carried out the reaction without any W, X or Y. Then she repeated the reaction three times adding a small amount of W, X or Y. In each case she timed how long the reaction took to finish. The results are in the table. substance added time of reaction/s 277 none W 266 279 Х Υ 78 (a) How would the student know when the reaction had finished? (b) (i) State the effect of each substance on the speed (rate) of the reaction. W Χ..... Y[3] (ii) Which substance, W, X or Y, is the best catalyst for this reaction? (c) How could the student check the reliability of her results? [Total: 7]

5 A student investigated the temperature changes when two different solids, **C** and **D**, dissolved in water.

7

For Examiner's Use

Two experiments were carried out.

Experiment 1

Using a measuring cylinder, 25 cm³ of distilled water was poured into a polystyrene cup. The initial temperature of the water was measured.

Solid **C** was added to the water, the timer started and the mixture stirred with a thermometer. The temperature of the solution was measured every 30 seconds for three minutes.

(a) Use the thermometer diagrams in the table to record the temperatures.

time/s	thermometer diagram	temperature/°C
0	30 25 20	
30	30 -25 -20	
60	25 20	
90	- 30 - 25 - 20	
120	30 -25 -20	
150	30 25 20	
180	30 25 20	

[2]

Experiment 2

The polystyrene cup was emptied and rinsed with water. Experiment 1 was repeated using solid **D** instead of solid **C**. The temperature of the solution was measured every 30 seconds for three minutes as before.


(b) Use the thermometer diagrams in the table to record the temperatures.

time/s	thermometer diagram	temperature/°C
0	25 20 15	
30	25 20 15	
60	25 20 15	
90	25 20 15	
120	25 20 15	
150	25 20 15	
180		

For Examiner's Use (c) Plot the results for Experiments 1 and 2 on the grid and draw two smooth line graphs. Clearly label your graphs.

9

For Examiner's Use



Show clearly on the graph how you worked out your answer.

.....S

[2]

(e)	What type of change occurs when substance D dissolves in water?
(f)	Suggest the effect on the results if Experiment 1 was repeated using 50 cm ³ of distilled water.
(g)	Predict the temperature of the solution in Experiment 2 after 1 hour. Explain your answer.
	[2]
(h)	When carrying out the experiments what would be the advantage of taking the temperature readings every 15 seconds?
	[2]
	[Total: 20]

https://xtremepape.rs/

For Examiner's Use

0620/62/M/J/12

6 Two substances, E and F, were analysed. E was a solid compound and F was a solution of ethanoic acid. The tests on E and F, and some of the observations, are in the table. Complete the observations

For Examiner's Use

in the table. Do not write any conclusions in the table.

		tests	observations
tests on solid E			
(a) Appearance of solid E.		pearance of solid E .	green powder
(b) Solid E was heated in a test-tube.The gas given off was tested.		id E was heated in a test-tube.	black solid formed
		e gas given off was tested.	limewater turned milky
(c)	(i)	Solid E was added to dilute sulfuric acid.	effervescence and blue solution formed
		The solution was divided into two equal portions in test-tubes.	
	(ii)	Excess aqueous sodium hydroxide was added to the first portion of the solution.	pale blue precipitate formed
(iii)	Drops of aqueous ammonia were added to the second portion of the solution.	pale blue precipitate formed
		Excess aqueous ammonia was then added to the mixture.	precipitate dissolved to form a dark blue solution
test	s on	liquid F	
(d) Appearance and smell of liquid F .		bearance and smell of liquid F .	appearance
(e) pH indicator paper was used to measure the pH of liquid F.			smell [2]
	the		pH[1]
((f)	Identify the gas given off in test (c)(i).	
			[1
((g)	Identify solid E .	
			[2
			[Total: 6

Fizzy drinks

12

© UCLES 2012

The bubbles in fizzy drinks are bubbles of carbon dioxide. The carbon dioxide is dissolved in

the drink under pressure. When a bottle of fizzy drink is opened the gas escapes and eventually the drink goes flat. The gas is lost more quickly if the fizzy drink is heated.

(a) How could the acidity of the fizzy drink be checked?

(b) Plan an experiment to find the volume of gas in a bottle of fizzy drink. You may use common laboratory apparatus.

......[6]

[Total: 7]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

For

Use