

Cambridge IGCSE[™]

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 0620/62

Paper 6 Alternative to Practical

May/June 2022

1 hour

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].


This document has 12 pages. Any blank pages are indicated.

IB22 06_0620_62/3RP © UCLES 2022

[Turn over

1 Sodium hydrogencarbonate decomposes when heated. The products are solid sodium carbonate, water and carbon dioxide.

A student decomposed a sample of sodium hydrogencarbonate using the apparatus shown.



(a)	Name the items of apparatus labelled A and B .
	A
	В
	[2]
(b)	When the sodium hydrogencarbonate was heated, a colourless liquid collected at the point marked ${\bf X}$.
	Suggest the identity of the colourless liquid.
	[1]
(c)	On the diagram draw one arrow to show where the apparatus should be heated during the experiment.
(d)	State an observation that would indicate the sodium hydrogencarbonate had stopped reacting.
	[1]
(e)	Explain why it is important to remove the delivery tube from the water as soon as heating is stopped.
	[2]


[Total: 7]

2 A student investigated the reaction between two different solutions of aqueous sodium carbonate, solution **K** and solution **L**, and dilute hydrochloric acid using two different indicators.

Two experiments were done.

Experiment 1

- A burette was rinsed with water and then with the dilute hydrochloric acid.
- The burette was filled with dilute hydrochloric acid. Some of the dilute hydrochloric acid was run out of the burette so that the level of the dilute hydrochloric acid was on the burette scale.
- Using a measuring cylinder, 25 cm³ of solution **K** was poured into a conical flask.
- Five drops of methyl orange indicator **and** five drops of thymolphthalein indicator were added to the conical flask.
- The conical flask was placed on a white tile.
- Dilute hydrochloric acid was added slowly from the burette to the conical flask, while the flask was swirled, until the solution turned yellow. This is the first colour change.
- More dilute hydrochloric acid from the burette was added to the conical flask, while swirling the flask, until the solution changed colour again. This is the second colour change.
- (a) Use the burette diagrams to complete the table for Experiment 1.




	Experiment 1
burette reading at first colour change/cm³	
final burette reading at second colour change/cm ³	
initial burette reading/cm³	
volume of dilute hydrochloric acid added for first colour change/cm ³	
total volume of dilute hydrochloric acid added for second colour change/cm³	

[3]

(b) Experiment 2

- The conical flask was emptied and rinsed with distilled water.
- Experiment 1 was repeated using solution L instead of solution K.

Use the burette diagrams to complete the table for Experiment 2.



initial burette reading

burette reading at first colour change

burette reading at second colour change

	Experiment 2
burette reading at first colour change/cm ³	
final burette reading at second colour change/cm ³	
initial burette reading/cm³	
volume of dilute hydrochloric acid added for first colour change/cm ³	
total volume of dilute hydrochloric acid added for second colour change/cm ³	

[3]

(c)	State the colour change observed at the end-point when dilute hydrochloric acid is added to	tc
	methyl orange in an alkaline solution.	

(d)	For Experiment 1, compare the volume of dilute hydrochloric acid needed for the first colour
	change with the volume of dilute hydrochloric acid for the second colour change.

[2]

© UCLES 2022 0620/62/M/J/22

(e)	e) Compare the concentration of solution K used in Experiment 1 to the concentration used in Experiment 2. Explain your answer.						
		[3					
(f)	(i)	Deduce the volume of dilute hydrochloric acid needed for the second colour change whe Experiment 2 is repeated using $50\mathrm{cm^3}$ of solution L .	n				
	(ii)	State why using 50 cm³ of solution L would cause a problem.	2]				
			 1]				
(g)	Sta	te the advantage of using a pipette instead of the measuring cylinder in these experiments					
(h)		plain why the conical flask was swirled as the dilute hydrochloric acid was added from th ette.	е				
		[
(i)	hyc	the start of Experiment 1, the burette was rinsed with water and then with dilut lrochloric acid. he start of Experiment 2, the conical flask was rinsed with water but not with solution L .	е				
	(i)	Explain why the conical flask was rinsed with water.					
		[
	(ii)	Explain why the conical flask was not rinsed with solution L in Experiment 2.					
		[
		[Total: 19	}]				

3 Solid **M** and solid **N** were analysed. Solid **M** was iron(III) nitrate. Tests were done on each substance.

tests on solid M

Complete the expected observations.

Solid \mathbf{M} was dissolved in water to form solution \mathbf{M} . Solution \mathbf{M} was divided into two approximately equal portions in two test-tubes.

(a)		the first portion of solution \mathbf{M} , aqueous sodium hydroxide was added gradually until in ess. The product was kept for (\mathbf{b}) .
	obs	ervations[2]
(b)	(i)	The product from (a) was transferred to a boiling tube. A piece of aluminium foil was added and the mixture warmed gently. Any gas produced was tested.
	(ii)	observations
	(,	[1]
(c)		the second portion of solution ${f M}$, about 1 cm depth of dilute nitric acid followed by a few ps of aqueous barium nitrate were added.
	obs	ervations[1]

tests on solid N

tests	observations
test 1	
A flame test was carried out on solid N .	the flame became red
Solid N was dissolved in water to form solution N . Solution N was divided equally into one test-tube and one boiling tube.	
test 2	
About 1 cm depth of dilute nitric acid followed by a few drops of aqueous silver nitrate were added to the first portion of solution N in a test-tube.	no visible change
test 3	
About 2cm depth of dilute hydrochloric acid was added to the second portion of solution N . The mixture was warmed and any gas produced was tested.	acidified aqueous potassium manganate(VII) changed from purple to colourless
(d) Identify the gas produced in test 3.	
	[1]
(e) Identify solid N.	
	[2]

[Total: 8]

4 The diagram shows some coffee beans.



Caffeine occurs naturally in coffee beans. Caffeine is a white crystalline solid. It is very soluble in hot water but much less soluble in cold water.

Plan an investigation to obtain a pure crystalline sample of caffeine from coffee beans.

Assume that all other soluble substances in coffee beans are very soluble in both hot and cold water.

You are provided with coffee beans and common laboratory apparatus.		
16.		

© UCLES 2022 0620/62/M/J/22

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.

© UCLES 2022 0620/62/M/J/22