

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

	CANDIDATE NAME			
	CENTRE NUMBER		CANDIDATE NUMBER	
*				
7 2	CHEMISTRY		0620/06	
2 2	Paper 6 Alternative to Practical		October/November 2007	
3			1 hour	
¢ 8	Candidates ans	wer on the Question Paper.		
3 7	No additional materials are required.			
2				

### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Exam	iner's Use
1	
2	
3	
4	
5	
6	
7	
Total	


This document consists of 13 printed pages and 3 blank pages.



- (a) Indicate on the diagram using arrows (i) where the copper sulphate crystals are placed, (ii) where heat is applied. [2] (b) What is the purpose of the ice? [1] (c) The crystals changed colour from \_\_\_\_\_\_to \_\_\_\_\_[2]
- **1** Hydrated copper sulphate crystals, CuSO<sub>4</sub>.5H<sub>2</sub>O were heated in the apparatus shown below.

[Total: 5]

2 An experiment was set up to investigate the rusting of iron.



3

**3** The information in the box is about the preparation of zinc nitrate crystals.

St	ep 1:	Add a small amount of zinc oxide to some hot dilute nitric acid, and stir.		
Step 2: Keep adding zinc oxide until it is in <i>excess</i> .				
Step 3: Remove the excess zinc oxide to leave colourless zinc nitrate solution.				
St	Step 4: Evaporate the zinc nitrate solution until it is saturated.			
St	ep 5:	Leave the saturated solution to cool. White crystals form on cooling.		
St	ep 6:	Remove the crystals from the remaining solution.		
St	ep 7:	Dry the crystals on a piece of filter paper.		
(a)	Sug	ggest a reason for using <i>excess</i> zinc oxide in Step 2.		
			[1]	
(b)	Sug	gest how the <i>excess</i> zinc oxide can be removed from the solution in Step 3.	[1]	
(c)	(i)	What is meant by the term saturated solution?		
	(ii)	What practical method could show the solution to be saturated?	[2]	
			[1]	
(d)	Why	are the crystals dried in Step 7 using filter paper instead of by heating?		
			[1]	

[Total: 6]

**4** A student investigated the reaction of dilute hydrochloric acid with two different solids, calcium carbonate (marble) and calcium oxide. Four experiments were carried out.

### Experiment 1

By using a measuring cylinder, 50 cm<sup>3</sup> of dilute hydrochloric acid was poured into a polystyrene cup and the initial temperature of the acid was measured. 2.5 g of small marble chips were added to the cup and the mixture stirred with the thermometer.

The temperature of the mixture was measured after 2 minutes.

Use the thermometer diagrams to record the temperatures in the table of results on page 6.





initial temperature/°C

final temperature/°C

#### Experiment 2

Experiment 1 was repeated using 2.5g of powdered calcium carbonate. Use the thermometer diagrams to record the results in the table.





initial temperature/°C

final temperature/°C

Experiment 3

Experiment 1 was repeated using 1.5 g of lumps of calcium oxide. Use the thermometer diagrams to record the temperatures in the table.



П	30
пП	25
	20

initial temperature/°C

final temperature/°C

### Experiment 4

Experiment 1 was repeated using 1.5g of powdered calcium oxide. Use the thermometer diagrams to record the results in the table.

6





initial temperature/°C


final temperature/°C

## Table of results

Experiment	temperature/°C		
Experiment	initial	final	difference
1			
2			
3			
4			

(a) What would be observed in Experiment 2?

[1]



7

(e) In Experiment 1, how would you know which reactant is in *excess*? Explain your answer.
[2]
(f) Explain how the temperature changes would differ in the experiments if 100 cm<sup>3</sup> of hydrochloric acid were used.
[2]
[1] [2]

5 Three different liquids **P**, **Q** and **R** were analysed. **Q** was an aqueous solution of sodium hydroxide.

The tests on the liquids and some of the observations are in the following table. Complete the observations in the table.


tests	observations	
<ul><li>(a) Test the pH of the liquids using indicator paper. Note the colour of the paper.</li></ul>	P colour red	
	Q colour[2]	
	R colour orange	
	рН 5	
(b) (i) Add a 5 cm piece of magnesium to about 3 cm <sup>3</sup> of liquid <b>P</b> in a test–tube. Test the gas given off.	bubbles of gas lighted splint pops	
(ii) Repeat (b)(i) using liquids <b>Q</b> , and <b>R</b> . Do not test for any gases.	Q R[2]	

For Examiner's Use

	-	-	
	tests	observations	
(c)	To about 2 cm <sup>3</sup> of liquid <b>P</b> add 1 spatula measure of sodium carbonate. Test the gas given off.		[3]
(d)	By using a teat pipette add aqueous silver nitrate to about 1 cm <sup>3</sup> of liquid <b>P</b> .	white precipitate	
(e)	By using a teat pipette add liquid <b>Q</b> to about 1 cm <sup>3</sup> of aqueous iron(II) sulphate.		[2]
( <b>f)</b> Name t	he gas given off in test <b>(b)(i)</b> .		[1]
( <b>g)</b> Name t	he gas given off in test <b>(c)</b> .		[1]
( <b>h)</b> Identify	liquid <b>P</b> .		[1]
(i) What co	onclusions can you draw about liqu	id <b>R</b> ?	
			 [2]

[Total: 14]

**6** Magnesium reacts with dilute sulphuric acid to form hydrogen gas. The speed of the reaction was investigated using the apparatus below.




In an experiment 50 cm<sup>3</sup> of dilute sulphuric acid was added to a large piece of magnesium. A student measured the total volume of gas produced at 2 minute intervals.

Use the measuring cylinder diagrams to complete the table.

time/minutes	measuring cylinder diagram	total volume of collected / cm <sup>3</sup>
0	5 10	
2	10 - 15 - 20	
4	25 	
6	25 - 30 - 35	
8		
10	40 45 50	
12	40 45 50	


[3]



(a) Plot the student's results on the grid. Use the points to draw a smooth line graph.

For Examiner's Use 7 Diesel is a liquid fuel obtained from crude oil. Biodiesel is a fuel made from oil obtained from the seeds of plants such as sunflowers.

Using the apparatus below plan an experiment to investigate which of these two fuels produces more energy.



[6]

[Total: 6]

# **BLANK PAGE**

# **BLANK PAGE**

### **BLANK PAGE**

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.