



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/32
Paper 3 (Extend	ded)	Oct	ober/November 2009
			1 hour 15 minutes

__._._

No Additional Materials are required.

Candidates answer on the Question Paper.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part questions.

For Examiner's Use		
1		
2		
3		
4		
5		
6		
7		
Total		

This document consists of 14 printed pages and 2 blank pages.



(a)	The	ne major gases in unpolluted air are 79% nitrogen and 20% oxygen.		
	(i)	Name another gaseous element in unpolluted air.		
	(ii)	Name two compounds in unpolluted air.		
		[2]		
(b)	Two	common pollutants in air are sulfur dioxide and the oxides of nitrogen.		
	(i)	Name another pollutant in air.		
	(ii)	Describe how sulfur dioxide is formed. [1]		
		[2]		
((iii)	How are the oxides of nitrogen formed?		
		ro1		
		[2]		
(c)	Hov	v is oxygen obtained from air?		
	•••••			
		[2]		
		[Total: 10]		

1

For Examiner's Use

[Total: 9]


(a)	(a) An important ore of zinc is zinc blende, ZnS.					
	(i) How is zinc blende changed into zinc oxide?					
			[1]			
	(ii)	Write a balanced equation for the reduction of zinc oxide to zinc by carbon.				
			[2]			
(b)		najor use of zinc is galvanizing; steel objects are coated with a thin layer of zinc. s protects the steel from rusting even when the layer of zinc is broken.				
		thin layer steel exposed to				
		of zinc oxygen and water				
		/ /3				
		steel				
	Explain, by mentioning ions and electrons, why the exposed steel does not rust.					
			[3]			

3

(c) Zinc electrodes have been used in cells for many years, one of the first was the Daniel cell in 1831.

For Examiner's Use

[Total: 10]



(i)	Give an explanation for the following in terms of atoms and ions.	
	observation at zinc electrode – the electrode becomes smaller	
	explanation	
		[1]
	observation at copper electrode – the electrode becomes bigger	
	explanation	
		[1]
(ii)	When a current flows, charged particles move around the circuit.	
	What type of particle moves through the electrolytes?	
		[1]
	Which particle moves through the wires and the voltmeter?	
		[1]

The distinctive smell of the seaside was thought to be caused by ozone, O₃. Ozone is a form of the element oxygen. (a) A mixture of oxygen and ozone is formed by passing electric sparks through oxygen. $3O_2 \rightleftharpoons 2O_3$ Suggest a technique that might separate this mixture. Explain why this method separates the two forms of oxygen. technique explanation [2] **(b)** Ozone is an oxidant. It can oxidise an iodide to iodine. $2I^{-} + O_{3} + 2H^{+} \rightarrow I_{2} + O_{2} + H_{2}O$ What would you see when ozone is bubbled through aqueous acidified potassium (i) iodide? (ii) Explain in terms of electron transfer why the change from iodide ions to iodine molecules is oxidation. (iii) Explain, using your answer to **b(ii)**, why ozone is the oxidant in this reaction.

For Examiner's Use

© UCLES 2009 0620/32/O/N/09

(c)		It is now known that the smell of the seaside is due to the chemical dimethyl sulfide, $(\text{CH}_3)_2\text{S}$.			
	(i)	Draw a diagram that shows the arrangement of the valency electrons in one molecule of this covalent compound. Use x to represent an electron from a carbon atom. Use o to represent an electron from a hydrogen atom. Use • to represent an electron from a sulfur atom.			
	(ii)	Name the three compounds formed when dimethyl sulfide is burnt in excess oxygen.	[3]		
			[2]		

[Total: 11]

For Examiner's Use

		t three elements in Group IV are carbon, silicon and germanium. ments and their compounds have similar properties.		
(a) The compound, silicon carbide, has a macromolecular structure similar to that of diamond.				
	(i)	A major use of silicon carbide is to reinforce aluminium alloys which are used in the construction of spacecraft. Suggest three of its physical properties.		
		TO.		
		[3]		
	(ii)	Draw a diagram to show the arrangement of silicon atoms around one carbon atom in silicon carbide. Label this diagram 1.		
		Draw a diagram to show the arrangement of carbon atoms around one silicon atom in silicon carbide. Label this diagram 2.		
		[3]		
(b) Germanium(IV) oxide, GeO_2 , has the same macromolecular structure as silicon(IV) oxide. Draw the structural formula of germanium(IV) oxide.				
		[2]		

© UCLES 2009 0620/32/O/N/09

5

:)	Germar	nium forms a series of hydrides comparable to the alkanes.	For Examiner's
	(i)	Draw the structural formula of the hydride which contains three germanium atoms per molecule.	Use
	(ii)	[1] Predict the products of the complete combustion of this hydride.	
	()	[2]	
		[Total: 11]	

6 (a) Sulfuric acid is made by the Contact process.

For
Examiner's
Use

2SO ₂	+	O_2	\rightleftharpoons	2SO ₃

(i)	Sulfur dioxide is made by burning sulfur. Name a source of sulfur.	
		••••
		[1]
(ii)	Give another use of sulfur dioxide.	
		[1]
(iii)	Name the catalyst used.	
		[1]
(iv)	If the temperature is decreased to 300 °C, the yield of sulfur trioxide increases. Explain why this lower temperature is not used.	
		[1]
(v)	Sulfur trioxide is dissolved in concentrated sulfuric acid. This is added to water to make more sulfuric acid. Why is sulfur trioxide not added directly to water?)
		[1]

© UCLES 2009 0620/32/O/N/09

(b)	Sulfuric acid was first made in the Middle East by heating the mineral, green vitriol, $FeSO_4.7H_2O$. The gases formed were cooled.									
	$\begin{array}{cccccccccccccccccccccccccccccccccccc$									
	$2FeSO_4(s) \rightarrow Fe2O_3(s) + SO_2(g) + SO_3(g)$									
	On cooling									
	$SO_3 + H_2O \rightarrow H_2SO_4$ sulfuric acid $SO_2 + H_2O \rightarrow H_2SO_3$ sulfurous acid									
	(i) How could you show that the first reaction is reversible?									
		[2]								
(ii) Sulfurous acid is a reductant. What would you see when acidified potassium manganate(VII) is added to a solution containing this acid?										
(iii) Suggest an explanation why sulfurous acid in contact with air changes into seacid.										
									[1]	
(c)	(c) 12.16 g of anhydrous iron(II) sulfate was heated. Calculate the mass of iron(III) oxide formed and the volume of gases, at r.t.p., formed.									
	$2FeSO_4(s) \ \to \ Fe_2O_3(s) \ + \ SO_2(g) \ + \ SO_3(g)$									
	mass of one mole of $FeSO_4 = 152 g$									
	number of moles of FeSO ₄ used =									
	number of moles of Fe_2O_3 formed =									
	mass of one mole of Fe_2O_3 =g									
	mass of iron(III) oxide formed =g									
	total number of moles of gases formed =									
	tota	I volume of gases formed =dm ³								

[6]

For Examiner's Use

[Total: 16]

7	Butan-1-ol is used as a solvent for paints and varnishes, to make esters and as a fuel. Butan-1-ol can be manufactured from but-1-ene, which is made from petroleum.										
	Biobutanol is a fuel of the future. It can be made by the fermentation of almost any form of biomass - grain, straw, leaves etc.										
	(a) But	(a) But-1-ene can be obtained from alkanes such as nonane, C ₉ H ₂₀ , by cracking.									
	(i)	Give the reaction conditions.									
			[2]								
	(ii)	Complete an equation for the cracking of nonane, C ₉ H ₂₀ , to give but-1-ene.									
		$C_9H_{20} \rightarrow$	[2]								
	(iii)	Name the reagent that reacts with but-1-ene to form butan-1-ol.									
			[1]								
	(b) (i)	Balance the equation for the complete combustion of butan-1-ol.									
		$\begin{array}{cccccccccccccccccccccccccccccccccccc$	[2]								
	(ii)	Write a word equation for the preparation of the ester butyl propanoate.									
			[2]								

(c)	The fermentation of biomass by bacteria produces a mixture of products which include biobutanol, propanol, hydrogen and propanoic acid.						
	(i)						
		propanol					
		propanoic acid					
		[2]					
	(ii)	Why is it important to develop these fuels, such as biobutanol, as alternatives to petroleum?					
		[1]					
(d)		v could you show that butanol made from petroleum and biobutanol are the same mical?					
		[1]					
		[Total: 13]					

© UCLES 2009 0620/32/O/N/09

BLANK PAGE

BLANK PAGE

DATA SHEET
The Periodic Table of the Elements

	0	4 He Heium	Neon 10 40	Argon	8 7	Krypton 36	131	Xenon 54	뜐	Radon 86		175 Lu Lutetium 71		۲	Lawrencium 103
	II/		19 Fluorine 9 35.5	Chlorine 17	≋ ಹ	0	127	lodine 53	Ą	Astatine 85		Yb Ytterbium 70		8	Nobelium 102
	VI		Oxygen 8		% Se	Selenium 34	128 7	Tellurium 52	Ро	_		169 Tm Thulium 69		Md	Mendelevium 101
	^		14 Nitrogen 7	Phosphorus 15	75 As	Arsenic 33	122	Antimony 51	508 B	Bismuth 83		167 Er Erbium 68			Fermium 100
	N		Carbon 6 Carbon 8 28	Silicon 14	_∞ 9	Germanium 32	£ 5		207 Pb	Lead 82		165 Ho Holmium 67			Einsteinium 99
	=		11 Baron 5 27 A1	Aluminium 13	° 8 B	Gallium 31	115	Indium 49	204 T 1	_ 18		162 Dy Dysprosium 66		ర	Californium 98
					₆₅ Zη	Zinc 30	115	Cadmium 48	201 Hg	Mercury 80		Tb Terbium 65			Berkelium 97
					²⁵ Ω	Copper 29		Silver 47	197 Au	Gold 79		157 Gd Gadolinium 64		Cm	Curium 96
Group					6 Έ	Nickel 28	901	Palladium 46	195 T	Platinum 78		152 Eu Europium 63		Am	Americium 95
Gro					ී දි	Cobalt 27	50 2	Rhodium 45	192 Ir	Iridium 77		Sm Samarium 62			Plutonium 94
		T Hydrogen			_{2е}	Iron 26	ē 2	Ruthenium 44	190 O S	Osmium 76		Pm Promethium 61		δ	Neptunium 93
					M Mn	Manganese 25	۲	1, 43	186 B	Rhenium 75		Neodymium 60	238		Uranium 92
					జ రే	Chromium 24	96 2	Molybdenum 42	² ≥	Tungsten 74		Pr Praseodymium 59		Ра	Protactinium 91
					15 >	Vanadium 23	88	-	18 E	Tantalum 73		140 Ce Cerium	232	드	Thorium 90
					48 F	Titanium 22	ار 9	Zirconium 40	178 ‡	72			nic mass	lod	nic) number
					Sc Sc	Scandium 21	8 >	_	139 La	Lanthanum 57 *	227 AC Actinium 89	series series	a = relative atomic mass	X = atomic symbol	b = proton (atomic) number
	=		9 Beryllium 4 Beryllium 24	Magnesium 12	⁶ С	Calcium 20	® ử	Strontium 38	137 Ba	Barium 56	226 Rad Radium 88	*58-71 Lanthanoid series 190-103 Actinoid series		×	Ω
	_		Lithium 3 Lithium 23 23	Sodium 11	≋ ⊻	Potassium 19	85	_	133 Cs	Caesium 55	Fr Francium 87	*58-71 L		Key	q

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.