

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

	CANDIDATE NAME						
	CENTRE NUMBER	CANDIE NUMBE					
8 8	CHEMISTRY					0	620/63
5 3 0	Paper 6 Alternat	tive to Practical	Oc	tobe	r/Nove		r 2013
4 3							1 hour
8 5	Candidates ans	wer on the Question Paper.					
2 3	No Additional M	laterials are required.					

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 9 printed pages and 3 blank pages.



1 A student tried to separate a mixture of salt and sand. Salt, sodium chloride, is soluble in water. Sand, silicon(IV) oxide, is insoluble in water. He added the mixture to water. Three sets of apparatus are shown.


For

Examiner's

Use

 $\times \times \times \times$ Α В С (a) Complete the boxes to identify the pieces of apparatus. [2] (b) Fill in the blanks in the following sentences. Liquids that dissolve substances are called The clear liquid formed is a [2] (c) Which set of apparatus should be used to obtain the sand, [2] salt crystals? (d) What happened to the water used in the experiment?[1] [Total: 7] 2 Sulfur dioxide is a poisonous gas which is denser than air and soluble in water. Sulfur dioxide can be prepared by adding dilute hydrochloric acid to sodium sulfite and warming the mixture.

Study the diagram of the apparatus used.



(a) Identify and explain three mistakes in the diagram.

	1	mistake
		reason
	2	mistake
		reason
	3	mistake
		reason
(b)	St	ate one precaution that should be taken when carrying out this experiment.
		[1]
		[Total: 7]

© UCLES 2013


For Examiner's Use

- **3** A scientist measured the boiling point of water at different pressures.
 - (a) Use the thermometer diagrams in the table to complete the boiling point temperatures.

pressure / atmospheres	thermometer diagram	boiling point /°C
1	105 100 	
2	125 120 	
3	140 135 130	
4	140 135 130	
5	155 - 150 - 145	
6	160 155 150	
7	170 165 160	

[3]

For Examiner's Use



4 A student investigated the reaction between aqueous potassium manganate(VII), which is purple, and two different colourless acidic solutions, **D** and **E**.

6


Three experiments were carried out.

(a) Experiment 1

A burette was filled with the solution of potassium manganate(VII) to the 0.0 cm^3 mark. Using a measuring cylinder, 25 cm^3 of solution **D** was poured into a conical flask.

Potassium manganate(VII) solution was added to the flask until the mixture just turned permanently pink.

Use the burette diagram to record the final volume in the table and complete the table.



final reading


	burette reading
final burette reading/cm ³	
initial burette reading/cm ³	
difference / cm ³	

[2]

(b) Experiment 2

Experiment 1 was repeated using 25 cm^3 of solution **E** instead of solution **D**.

Use the burette diagrams to record the readings in the table and complete the table.



initial reading

final reading

	burette reading
final burette reading/cm ³	
initial burette reading/cm ³	
difference/cm ³	

For

Examiner's

Use

(c)	Exp	periment 3	For Examiner's
	obs	eous ammonia was added to solution E in a test-tube. A green precipitate was erved. mixture was left to stand for 5 minutes. The surface of the precipitate turned brown.	Use
	Wha	at conclusions can you draw from these observations?	
		[3]	
(d)	(i)	What colour change was observed as potassium manganate(VII) solution was added to the flask in Experiment 1?	
		[1]	
	(ii)	Why was an indicator not added to the flask?	
	()		
		[1]	
(e)	(i)	In which experiment was the greatest volume of potassium manganate(VII) solution used?	
	(ii)	Compare the volumes of potassium manganate(VII) used in Experiments 1 and 2.	
		[1]	
	(iii)	Suggest an explanation for the difference in volumes.	
		[2]	
(f)		xperiment 2 was repeated using 12.5 cm ³ of solution E , what volume of potassium nganate(VII) solution would be used? Explain your answer.	
(g)	Give and	e one advantage and one disadvantage of using a measuring cylinder for solutions D E .	
	adv	antage	
	disa	advantage[2]	
		[Total: 17]	

5 Two liquids, **F** and **G**, were analysed. **G** was an aqueous solution of potassium chloride. The tests on the liquids and some of the observations are in the following table. Complete the observations in the table.

For Examiner's Use

	tests	observations		
tests or	tests on liquid F			
(a) (i) Appearance of liquid F.		yellow solution		
	The pH of the liquid was tested.	pH = 7		
(ii) An equal volume of dilute sulfuric acid was added to liquid F.		solution turned orange		
	Excess aqueous sodium hydroxide was then added to the mixture.	solution turned from orange to yellow		
 (b) Dilute sulfuric acid was added to liquid F followed by hydrogen peroxide. 		rapid effervescence		
The mixture was shaken and the gas given off tested with a splint.		glowing splint relit		
tests on liquid G				
(c) Dilute nitric acid was added to liquid G followed by aqueous barium nitrate.				
. ,	ute nitric acid was added to liquid G lowed by aqueous silver nitrate.	[2]		
(e) What does test (a)(i) tell you about liquid F?				
(f)	(f) What type of reaction happened in test (a)(ii)? Explain your answer.			
type of reaction				
	explanation			
(g) Identify the gas given off in test (b).				
		[Total: 8]		

6

Indigestion tablets

Indigestion pain is caused by too much acid in your stomach. The acid is hydrochloric acid. Indigestion tablets contain a base which neutralises the acid.

You are provided with two different brands of indigestion tablets, Painremuve and Indcure.

Plan an investigation to compare which of these brands of tablet is the most effective. You are provided with dilute hydrochloric acid and common laboratory apparatus.

[Total: 7]

For

Examiner's

Use

9

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

BLANK PAGE