

Cambridge Assessment International Education Cambridge International General Certificate of Secondary Education

CHEMISTRY

0620/43 October/November 2018

Paper 4 Extended Theory MARK SCHEME Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2018 series for most Cambridge IGCSE[™], Cambridge International A and AS Level components and some Cambridge O Level components.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **11** printed pages.

[Turn over

© UCLES 2018

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a guestion. Each guestion paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

the specific content of the mark scheme or the generic level descriptors for the question the specific skills defined in the mark scheme or in the generic level descriptors for the question the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded positively:

marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate

marks are awarded when candidates clearly demonstrate what they know and can do

marks are not deducted for errors

marks are not deducted for omissions

answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the guestion as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

| Question | Answer | Marks |
|----------|-----------------|-------|
| 1(a) | oxygen | 1 |
| 1(b) | hematite | 1 |
| 1(c) | sulfur dioxide | 1 |
| 1(d) | ammonia | 1 |
| 1(e) | carbon monoxide | 1 |
| 1(f) | sodium chloride | 1 |
| 1(g) | carbon dioxide | 1 |
| 1(h) | oxygen | 1 |

| Question | Answer | Marks |
|----------|--|-------|
| 2(a)(i) | M1 breakdown of an ionic compound when molten or in aqueous solution | 2 |
| | M2 (using) electricity / electric current | |
| 2(a)(ii) | M1 electron(s) | 2 |
| | M2 ion(s) | |
| 2(b)(i) | M1 inert / unreactive | 2 |
| | M2 conducts electricity | |

0620/43

| Question | | An | swer | | Marks |
|----------|---------------------------|-----------------------------|------------------------------|-------------------------------|-------|
| 2(b)(ii) | observation at anode(+) | name of product at anode(+) | observation at cathode(–) | name of product at cathode(-) | 6 |
| | M1 green / yellow bubbles | M2 chlorine | | M3 hydrogen | |
| | | M4 oxygen | M5 pink / brown solid | M6 copper | |

| Question | Answer | Marks |
|----------|--|-------|
| 3(a) | $[(64 \cdot 2) + 56 + 119 + (32 \cdot 4) =] 431$ | 1 |
| 3(b) | [(119/151) · 100 =] 78.8 (%) | 1 |
| 3(c) | SnO ₂ because the percentage of tin is larger in SnO ₂ or answer to (b) > 27.6 % | 1 |
| 3(d) | $SnO_2 + 2C \rightarrow Sn + 2CO$ | 2 |
| | M1 all formulae correct | |
| | M2 equation fully correct | |
| 3(e) | M1 (\rightarrow) Fe ²⁺ + Sn OR 2Fe + 3Sn ²⁺ \rightarrow 2Fe ³⁺ + 3Sn | 2 |
| | $ \begin{array}{l} \textbf{M2} (\rightarrow) \ \textbf{Sn}^{2+} + \textbf{Cu} \\ \textbf{OR} \ \textbf{Sn} + 2 \textbf{Cu}^{2+} \rightarrow \textbf{Sn}^{4+} + 2 \textbf{Cu} \end{array} $ | |
| 3(f)(i) | M1 glowing splint | 2 |
| | M2 relights / rekindles | |

| | | FODLIGHED | | | |
|-----------|--|---|---|--|--|
| Question | Answer | | | | |
| 3(f)(ii) | M1 nitrogen dioxide / nitrogen(IV) oxide | | | | |
| | M2 brown (gas) | | | | |
| 3(f)(iii) | $2Cu(NO_3)_2 \rightarrow 2CuO + 4NO_2 + O_2$ | | 1 | | |
| 3(g)(i) | zinc acts as a barrier which prevents contact between iron and water and air / oxygen | | | | |
| 3(g)(ii) | SUMMARY | | | | |
| | N | 1 comparison of reactivity | | | |
| | N | 2 zinc loses electrons | | | |
| | N | 3 where electrons move to OR iron does not lose electrons | | | |
| | M1 zinc is more reactive than iron / steel ORA | | | | |
| | M2 zinc loses electrons / zinc is oxidised | | | | |
| | M3 electrons are transferred to iron / iron is not oxidised / iron does not lose electrons | | | | |

| Question | Answer | Marks |
|----------|---|-------|
| 4(a) | M1 (Mol KOH =) $0.00125 / 1.25 \cdot 10^{-3}$ | 3 |
| | M2 (Mol H ₂ SO ₄ =) 0.000625 / $6.25 \cdot 10^{-4}$ | |
| | M3 (Conc H ₂ SO ₄ =) $0.03125 / 3.125 \cdot 10^{-2} (mol / dm3)$ | |

| Question | | | Answer | Marks |
|-----------|------------------------------------|-------------------|---|-------|
| 4(b) | SUMMARY | | | |
| | Γ | M1 | repeat | |
| | | M2 | heat (liquid or solution should be implied) | |
| | | М3 | when to stop heating | |
| | | M4 | what to do after heating | |
| | | M5 | method of drying crystals (crystals or solid should be implied) | |
| | M1 repeat without indic | ator us | sing same volumes | |
| | M2 evaporate / heat / wa | arm / b | ooil / leave in sun | |
| | M3 until most of the wa | iter is g | gone / some water left / saturation(point) / crystallisation point / evaporate some of the water | |
| | M4 leave / (allow to) cod | ol / allo | ow to crystallise | |
| | M5 details of drying | | | |
| 4(c)(i) | M1 bubbles / effervesce | ence / f | fizzing | |
| | M2 solid or magnesium | n disso | lves / solid or magnesium disappears | |
| 4(c)(ii) | lilac flame | | | |
| 4(c)(iii) | white precipitate | | | |
| 4(d)(i) | $Mg(OH)_2 + H_2SO_4 \rightarrow M$ | lgSO ₄ | + 2H ₂ O | 2 |
| | M1 formula of both Mg | (OH) ₂ | and MgSO₄ | |
| | M2 equation fully correct | ct | | |

https://xtremepape.rs/

| Question | Answer | Marks | | |
|-----------|---|-------|--|--|
| 4(d)(ii) | $Zn + H_2SO_4 \rightarrow ZnSO_4 + H_2$ | 2 | | |
| | M1 formula of ZnSO₄ | | | |
| | M2 equation fully correct | | | |
| 4(d)(iii) | $Na_2CO_3 + H_2SO_4 \rightarrow Na_2SO_4 + CO_2 + H_2O$ | 2 | | |
| | M1 formulae of both Na ₂ CO ₃ and Na ₂ SO ₄ | | | |
| | M2 equation fully correct | | | |

| Question | Answer | Marks |
|----------|--|-------|
| 5(a) | M1 volume of gas | 2 |
| | M2 time | |
| 5(b) | M1 rate decreases / reaction gets slower | 3 |
| | M2 concentration of acid decreases | |
| | M3 fewer collisions per unit time | |

| Question | Answer | Marks |
|----------|---|-------|
| 5(c) | M1 particles have more kinetic energy | 4 |
| | M2 particles move faster | |
| | M3 more collisions per unit time | |
| | M4 more of the particles have energy greater than or equal to activation energy / more of the collisions have energy greater than or equal to activation energy | |
| | OR more of the particles have sufficient energy to react / more of the collisions have sufficient energy to react | |
| | OR A great er percentage or great er proportion or great er fraction of collisions are successful | |
| 5(d) | ANY TWO FROM: increase concentration of hydrochloric acid decrease particle size of calcium carbonate / increase surface area of calcium carbonate (add)catalyst | 2 |

| | | PUBLISH | ED | | |
|-----------|---|-------------|-----------|-----------------------|-------|
| Question | | Answ | er | | Marks |
| 6(a)(i) | SUMMARY | | | | 6 |
| | M1 and | M4 re | eactants | | |
| | M2 and | M5 c | onditions | | |
| | M3 and | M6 e | quation |] | |
| | FERMENTATION: M1 glucose / sucrose / starch / other named carbohydra M2 Zymase / Yeast / 37° C M3 C ₆ H ₁₂ O ₆ \rightarrow 2C ₂ H ₅ OH + 2CO ₂ HYDRATION: M4 Ethene and steam or water can score in equation a M5 H ₃ PO ₄ (catalyst) / 300° C / 60 atm M6 C ₂ H ₄ + H ₂ O \rightarrow C ₂ H ₅ OH | | | on as correct formula | |
| 6(a)(ii) | ANY TWO FROM:- carbohydrates are renewable fossil fuels are non-renewable lower temperature means fossil fuels conserved C lower temperature means lower energy costs OR hydration reaches an equilibrium meaning lower y | Α | | | 2 |
| 6(a)(iii) | M1 solvent | | | | 2 |
| | M2 fuel | | | | |
| 6(b)(i) | E | | | | 1 |

| https:/ | 0620/43 | |
|---------------|-----------|---|
| //xtrem | Question | |
| iepap | 6(b)(ii) | D |
| 9. rs/ | 6(b)(iii) | В |
| | 6(b)(iv) | С |

| Question | Answer | Marks |
|-----------|--------|-------|
| 6(b)(ii) | D | 1 |
| 6(b)(iii) | В | 1 |
| 6(b)(iv) | C | 1 |
| 6(b)(v) | Α | 1 |