

Cambridge IGCSE™

CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/63 May/June 2022

Paper 6 (Extended) MARK SCHEME Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the May/June 2022 series for most Cambridge IGCSE, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

| Ma | Maths-Specific Marking Principles | | | | | |
|----|---|--|--|--|--|--|
| 1 | Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing. | | | | | |
| 2 | Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected. | | | | | |
| 3 | Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points. | | | | | |
| 4 | Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw). | | | | | |
| 5 | Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread. | | | | | |
| 6 | Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear. | | | | | |

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M Method marks, awarded for a valid method applied to the problem.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation '**dep**' is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

- awrt answers which round to
- cao correct answer only
- dep dependent
- FT follow through after error
- isw ignore subsequent working
- nfww not from wrong working
- oe or equivalent
- rot rounded or truncated
- SC Special Case
- soi seen or implied

| Question | Answer | Marks | Partial Marks |
|----------|---|-------|---------------|
| 1(a) | 3, 4, 5, 6 in correct rows | 1 | |
| 1(b) | R = n | 1 | |
| 2(a) | Circle with 3, 4 or 5 diameters drawn | C1 | |
| | 6, 8, 10 in correct rows | 1 | |
| 2(b) | R = 2n | 1 | |
| 3(a) | 11 | 1 | |
| 3(b) | 16 in row 5 | 1 | |
| 3(c) | Three second differences of 1 seen AND coefficient of n^2 equal to $\frac{1}{2}$ seen | C1 | |
| | or use of quadratic expression seen $\frac{2}{2}$ | | |
| | Derivation of coefficient of <i>n</i> equal to $\frac{1}{2}$ seen | C1 | |
| | Derivation of constant equal to 1 seen | C1 | |
| | $R = \frac{1}{2}n^2 + \frac{1}{2}n + 1$ | 1 | |
| | 3(c) alternative 1 | | |
| | Three second differences of 1 seen AND coefficient of n^2 equal to $\frac{1}{2}$ seen | C1 | |
| | $[\frac{1}{2}n^2]$ 0.5 2 4.5 8 12.5 | C1 | |
| | [leaves] 1.5 2 2.5 3 3.5 One difference of 0.5 seen (b =) 0.5 | C1 | |
| | $R = \frac{1}{2}n^2 + \frac{1}{2}n + 1$ | 1 | |

0607/63

| Question | Answer | Marks | Partial Marks |
|----------|---|-------|---------------|
| 3(c) | 3(c) alternative 2 | | |
| | Three second differences of 1 seen AND | C1 | |
| | coefficient of n^2 equal to $\frac{1}{2}$ seen | | |
| | $ \begin{array}{cccccccccccccccccccccccccccccccccccc$ | C1 | |
| | a+b=1 indicated | C1 | |
| | $R = \frac{1}{2}n^2 + \frac{1}{2}n + 1$ | 1 | |
| 4(a) | [tangents] do not intersect or [tangents are] parallel | 1 | |
| 4(b) | 10 | 1 | |
| 4(c) | 4th tangent drawn that intersects the other 3 tangents | 1 | |
| | 15 (on row 4 of table) | 1 | |
| 4(d) | Equation with correct substitution for R and n | C1 | |
| | 1.5 oe | 1 | |
| 5(a) | 3 secants intersecting each other | C1 | |
| | 13 | 1 | |
| 5(b) | Two equations with correct substitution for R and n | 1 | |
| | Correct method to eliminate either <i>b</i> or <i>c</i> | C1 | |
| | (b =) 2.5 (c =) 1 | 2 | B1 for each |
| | 5(b) alternative 1 | | |
| | $\begin{bmatrix} 1/_2 n^2 \end{bmatrix} \begin{array}{ccccccccccccccccccccccccccccccccccc$ | 1 | |
| | One difference of 2.5 seen | C1 | |
| | (b =) 2.5 (c =) 1 | 2 | B1 for each |

| Question | | | | Answ | | Marks | Partial Marks | | | |
|----------|---|----------|----------------------|-----------|---------|------------|---------------|----|----|---|
| 5(b) | 5(b) alternative 2 | | | | | | | | | |
| | 1 4 8 13 19 26 3 4 5 6 7 | | | | | | | | 1 | |
| | <i>a</i> + <i>b</i> = | 3 indica | ted | | | | | | C1 | |
| | (b =) 2 (c =) 1 | 5 | | | | | | | 2 | B1 for each |
| 6 | Sketch or corre | | adratic tution ar | nd rearra | ingemer | t to $= 0$ | | | C1 | FT their 4(d) |
| | Correct intersection of curve and straight line marked on sketch or correct factorisation or correct substitution into quadratic formula | | | | | | | | C1 | |
| | 48 | | | | | | | | 1 | |
| 7(a) | Their $5(b) - their 3(c) = 60$ oe Or list of results showing differences | | | | | | | | C1 | |
| | 30 | | | | | | | | 1 | |
| 7(b) | Correct substitution of <i>their</i> 30 into 5(b) or 3(c) | | | | | | | | C1 | FT their 5(b) or their 3(c) |
| | 526 | | | | | | | | 1 | |
| 8(a) | Correct calculation for a correct figure in table | | | | | | | | C1 | |
| | Plan | inter | arrive | start | land | end | wait | | 4 | B1 for each cell |
| | D | 25 | 535 | 610 | 280 | 890 | 75 | | | FT for 295, their 890 – their 595 |
| | E 60 595 890 44 934 295 | | | | | | | | | |
| 8(b) | 8(b) $1080 - (150 + 110 + 100 + 280 + 44)$ oe | | | | | | | C1 | | |
| | 396 | | | | | | | 1 | | |

| Question | | | Ansv | | Marks | Partial Marks | | |
|----------|--|--------------------------|-----------|-------------------|-------|---------------|---|---|
| 9(a) | Inter | Num | % | | Cum % | | 3 | B1 for 16 |
| | 60< <i>t</i> ≤ 120 | 34 | 19 | <i>t</i> ≤ 120 | 42 | | | B2 for other 6 cells correct |
| | 120< <i>t</i> ≤ 180 | 29 | 16 | $t \leqslant 180$ | 58 | | | Or B1 for 3, 4 or 5 other cells correct |
| | 180< <i>t</i> ≤ 240 | 23 | 13 | $t \leq 240$ | 71 | | | FT <i>their</i> 16 for 58 |
| | 240< <i>t</i> ≼ 300 | 16 | 9 | <i>t</i> ≤ 300 | 80 | | | FT <i>their</i> 58 for 13 |
| | 300< <i>t</i> ≼ 360 | 11 | 6 | <i>t</i> ≤ 360 | 86 | | | |
| | 360< <i>t</i> ≤420 | 11 | 6 | <i>t</i> ≤ 420 | 92 | | | |
| | 420< <i>t</i> ≤480 | 7 | 4 | $t \leqslant 480$ | 96 | | | |
| | 480< <i>t</i> ≤ 540 | 4 | 2 | $t \leq 540$ | 98 | | | |
| | 540< <i>t</i> ≼600 | 2 | 1 | $t \leqslant 600$ | 99 | | | |
| 9(b) | Points correct | ly plotte | d | | | | 1 | FT their points |
| | Correct curve | : | | | | | 1 | FT their plotted points |
| 9(c) | Between 140 | and 160 | | | | | 1 | |
| 10(a) | 23 = k(60 - | $a)^{\frac{1}{3}}$ isw | | | | | 1 | |
| | 23 = k (60 - k) $86 = k (360 - k)$ | $(-a)^{\frac{1}{3}}$ isw | 7 | | | | | |
| 10(b) | 86 (360- | $(a)^{\frac{1}{3}}$ | | | | | 1 | |
| | $\frac{86}{23} = \frac{(360-a)^{\frac{1}{3}}}{(60-a)^{\frac{1}{3}}}$ | | | | | | | |
| | $\left(\frac{86}{23}\right)^3 = \frac{360 - a}{60 - a}$ | | | | | | | |
| 10(c) | 52.3(60-a) | = 360 - | a | | 1 | | | |
| | Correctly isol | ating a | | | C1 | | | |
| | 54.1 [] or 5 | 4.2 leadi | ng to 54 | | 1 | | | |
| 10(d) | Correct substi | itution of | 54 into c | rt (a) | C1 | | | |
| | $p = \dots 13 \dots (t - \dots 54 \dots)^{\frac{1}{3}}$ | | | | | | 1 | FT |

| Question | Answer | Marks | Partial Marks |
|----------|--|-------|---|
| 10(e) | $50 = their 13(t-54)^{\frac{1}{3}}$ or correct sketch of model | 1 | |
| | Correct first step or correct sketch with line at 50 | C1 | |
| | 111 | 1 | FT their 13 |
| 10(f) | $\underbrace{Correct sketch}_{0}$ | 2 | FT <i>their</i> model from 10(d)B1 for correct shapeB1 for position |
| 10(g) | Valid up to $t = approx. 500$ or invalid after $t = approx. 500$ | 1 | FT <i>their</i> model |
| 11(a) | 52 to 53 | 1 | FT <i>their</i> model from 10(d) |
| 11(b) | 95 | 1 | |
| 11(c) | Yes, with appropriate comment e.g. approx. half the planes have to wait to land | 1 | |