



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/02

Paper 2 (Extended)

October/November 2013

45 minutes

Candidates answer on the Question Paper.

Additional Materials: Geometrical Instruments

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

Do not use staples, paper clips, highlighters, glue or correction fluid.

You may use a pencil for any diagrams or graphs.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

CALCULATORS MUST NOT BE USED IN THIS PAPER.

All answers should be given in their simplest form.

You must show all the relevant working to gain full marks and you will be given marks for correct methods even if your answer is incorrect.

The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 40.

This document consists of 8 printed pages.



Formula List

For the equation

$$ax^2 + bx + c = 0$$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Curved surface area, A, of cylinder of radius r, height h.

$$A = 2\pi rh$$

Curved surface area, A, of cone of radius r, sloping edge l.

$$A = \pi r l$$

Curved surface area, A, of sphere of radius r.

$$A = 4\pi r^2$$

Volume, V, of pyramid, base area A, height h.

$$V = \frac{1}{3}Ah$$

Volume, V, of cylinder of radius r, height h.


$$V = \pi r^2 h$$

Volume, V, of cone of radius r, height h.

$$V = \frac{1}{3}\pi r^2 h$$

Volume, V, of sphere of radius r.

$$V = \frac{4}{3}\pi r^3$$



$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cos A$$

Area =
$$\frac{1}{2}bc \sin A$$

Answer all the questions.


For Examiner's Use

1 Solve the simultaneous equations.

$$3g - 2h = 11$$
$$g - 2h = 5$$

$$h =$$
 [2]

2 (a)





$$n(U) = 20, \ n(A \cup B)' = 3, \ n(A) = 11, \ n(B) = 13.$$

Find $n(A \cap B')$.

Answer(a)	[2]
Answer(a)	12

(b) On each Venn diagram, shade the region indicated.





[2]

3 Tiago buys a concert ticket and then sells it for \$15. He makes a profit of 20%.

Calculate how much Tiago paid for the ticket.

Answer \$[3]

4

D36° В Q

Examiner's Use

ABCD is a parallelogram and BQPC is a rhombus.

DCE is a straight line. Angle $DAB = 52^{\circ}$ and angle $ECP = 36^{\circ}$.

Find the size of angle *BPC*.

Answer	[3]
--------	-----

NOT TO **SCALE**

(a) Simplify $\sqrt{72}$.

(b)
$$\frac{\sqrt{2}+2}{\sqrt{2}-1} = p+q\sqrt{2}$$

Find the values of p and q.

$$Answer(b) p =$$

$$q =$$
 [3]

© UCLES 2013

0607/02/O/N/13

For Examiner's Use

6	Sim	plify the following.					
	(a)	$2y^2 \times 3y^3$					
	(b)	Answer(a)[2] $\sqrt[3]{27p^{27}}$					
		Answer(b) [2]					
7	(a)	Find the amplitude and period of the function $f(x) = 4\cos(4x)$.					
		Answer(a) Amplitude =					
		Period =					
	(b)	$g(x) = 4\cos(4x) - 4$					
		Describe fully the single transformation that maps the graph of $y = f(x)$ onto the graph $y = g(x)$.					
		Answer(b)					
		[2]					


8 (a) Write down the value of $8^{\frac{1}{3}}$.

For Examiner's Use

- Answer(a) [1]
- **(b)** Find the exact value of $\left(\frac{4}{3}\right)^{-2}$.

Answer(b) [2]

9



NOT TO SCALE


The diagram shows a cube of side length 1.

Find the value of $\tan x^{\circ}$.

Answer [3]

© UCLES 2013 0607/02/O/N/13

10




For Examiner's Use

Find the exact value of x.

Answer
$$x =$$
 [3]

11



NOT TO SCALE

ABCD is a parallelogram. DM = MC and CN = 2NB. $\overrightarrow{AB} = \mathbf{p}$ and $\overrightarrow{AD} = \mathbf{q}$.

(a) Write down \overrightarrow{CN} in terms of q.

Answer(a) [1]

(b) Find \overrightarrow{MN} in terms of **p** and **q**.

Answer(b) [1]

Question 12 is printed on the next page.

		o	
12	f(x) = 3x - 1	g(x) = 12 - x	
	Find		
	(a) $f(g(8))$,		
		Answer(a)	
	(b) $f(g(x))$, in its simplest form,	Tins wer (a)	
		Answer(b)	
	(c) $g^{-1}(x)$.	(-)	

Answer(c) $g^{-1}(x) =$ [1]

Examiner's Use

[2]

[2]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

© UCLES 2013

0607/02/O/N/13