

## **Cambridge International Examinations**

Cambridge Ordinary Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

BIOLOGY 5090/62

Paper 6 Alternative to Practical

May/June 2017

1 hour

Candidates answer on the Question Paper.

No Additional Materials are required.

## **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Write your answers in the spaces provided on the Question Paper.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

This document consists of 8 printed pages.



## Answer **all** the questions in the spaces provided.

1 (a) Some students investigated the effect of different concentrations of sucrose solution on potato tissue.

Four strips of potato **A**, **B**, **C** and **D**, were cut. Each strip measured  $80 \,\text{mm} \times 4 \,\text{mm} \times 4 \,\text{mm}$ . The mass of each strip was measured and recorded in Table 1.1.

One strip of potato was placed in each of four sucrose solutions of different concentrations:

- 0.2 mol per dm<sup>3</sup>
- 0.4 mol per dm<sup>3</sup>
- 0.6 mol per dm<sup>3</sup>
- 0.8 mol per dm<sup>3</sup>

The same volume of sucrose solution was used for each strip.

The strips were left for 30 minutes. After 30 minutes, the strips were removed from the sucrose solutions and carefully blotted dry. The mass of each strip was then measured again and recorded in Table 1.1.

Table 1.1

potato strip	concentration of sucrose solution /mol per dm <sup>3</sup>	mass of potato strip at start /g	mass of potato strip after 30 minutes/g	change in mass /g
Α	0.2	4.0	4.3	
В	0.4	4.0	4.1	
С	0.6	4.0	3.8	
D	0.8	4.0	3.6	


(i)	Complete Table 1.1 by calculating the change in mass for each potato strip.	[3]
(ii)	Suggest explanations for the results for strip <b>A</b> and strip <b>D</b> .	
		[4]

© UCLES 2017 5090/62/M/J/17

(iii)	Suggest why each strip was blotted dry after being removed from the sucrose solution.
	[1]
(iv)	Explain why all the strips were cut to the same size ( $80\text{mm} \times 4\text{mm} \times 4\text{mm}$ ) at the start of this investigation.
	[2]
(v)	You are given 50 cm <sup>3</sup> of a sucrose solution containing 0.8 mol per dm <sup>3</sup> . Describe how you would use this solution to prepare 100 cm <sup>3</sup> of 0.4 mol per dm <sup>3</sup> sucrose solution.
	[1]

**(b)** When plant cells lose water, the cytoplasm may shrink and move away from the cell wall. When this happens, the cells are **plasmolysed**.

Fig. 1.1 represents a group of plant cells, some of which are plasmolysed.



key	
	plasmolysed cell
	non-plasmolysed cell

Fig. 1.1

(i) Complete Table 1.2 by counting the number of plasmolysed cells and the number of non-plasmolysed cells.

Table 1.2

number of plasmolysed cells	number of non-plasmolysed cells

[1]

(ii) Calculate the number of plasmolysed cells as a percentage of the total number of cells.
Show your working.

 	 	 	 	 	 				 		 								 	٠.	9	6
																				Γ	2	)

**(c)** A student carried out an investigation into the relationship between the concentration of sucrose solution and the number of plant cells which were plasmolysed.

She placed small pieces of plant tissue in sucrose solutions and counted the number of cells that were plasmolysed. She then calculated the percentage of cells that were plasmolysed in each solution.


Her results are shown in Table 1.3.

Table 1.3

concentration of sucrose solution/mol per dm <sup>3</sup>	percentage of cells that were plasmolysed
0.0	0
0.2	5
0.4	18
0.6	75
0.8	100

© UCLES 2017 5090/62/M/J/17

(i) Plot a line graph of the results in Table 1.3. Join the points on your graph with ruled, straight lines.



[4]

(ii) Use your graph to find the concentration of sucrose solution in which 50% of the cells would be plasmolysed. On your graph, show how you obtained this value.

Concentration of sucrose solution in which 50% of the cells would be plasmolysed:

1	2

[Total: 20]

ips 11.0 1050 4.0	(	(i) star	ch			
reducing sugars						
reducing sugars						
ble 2.1 gives information about the composition of some foods.  Table 2.1  food fat/g per 100 g energy/kJ per 100 g protein/g per 10  hips 11.0 1050 4.0  hicken 5.0 630 25.0  reet potato 0.6 360 1.0						
Table 2.1  food	(	ii) redu	icing suga	ars		
Table 2.1  food						
Table 2.1  food						
Table 2.1  food						
Table 2.1  food						
Table 2.1  food						
Table 2.1           food         fat/g per 100 g         energy/kJ per 100 g         protein/g per 10 g           nips         11.0         1050         4.0           hicken         5.0         630         25.0           veet potato         0.6         360         1.0						
food         fat/g per 100 g         energy/kJ per 100 g         protein/g per 10 g           nips         11.0         1050         4.0           hicken         5.0         630         25.0           veet potato         0.6         360         1.0						
hicken 5.0 630 25.0 veet potato 0.6 360 1.0	(b) <sup>-</sup>	  Table 2.1	gives inf			
hicken 5.0 630 25.0 veet potato 0.6 360 1.0	(b) <sup>-</sup>	  Table 2.1	gives inf	ormation about the co	mposition of some foods.	
veet potato 0.6 360 1.0	(b) -	Table 2.1	gives inf	ormation about the co	mposition of some foods.	
·			gives inf	ormation about the co Tat fat/g per 100 g	mposition of some foods.  ole 2.1  energy/kJ per 100 g	protein/g per 100 g
as 0.4 210 5.0	otato	food		ormation about the co  Tat  fat/g per 100 g  11.0	mposition of some foods.  ole 2.1  energy/kJ per 100 g  1050	protein/g per 100 g 4.0
	otato	food chips d chicker	1	ormation about the co  Tat  fat/g per 100 g  11.0  5.0	mposition of some foods.  Die 2.1  energy/kJ per 100 g  1050  630	protein/g per 100 g 4.0 25.0
Using the information in Table 2.1, state the relationship	otato	<b>food</b> chips		ormation about the co  Tat  fat/g per 100 g  11.0	mposition of some fole 2.1  energy/kJ per 1  1050	oods.

© UCLES 2017 5090/62/M/J/17

(ii)	Calculate the protein content of 250 g of cooked chicken.
	Show your working.
	[2
(iii)	Calculate the mass of boiled peas that you would need to eat to obtain the same mass of protein as in 100 g of cooked chicken.
	Show your working.
	[2
	[Total: 10


Question 3 starts on the next page.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

**3** Fig. 3.1 shows cells as seen using a light microscope.



## magnification ×200

Fig. 3.1

(a) In the space below, make a large drawing of the cell labelled **P**. You do not need to label your drawing.

(b)	Measure and record the maximum length of cell <b>P</b> in Fig. 3.1.	
	Maximum length of cell <b>P</b> in Fig. 3.1 mm	
	Use the magnification of Fig. 3.1 to calculate the <b>actual</b> length of cell <b>P</b> .	
	Show your working.	
		 [4]
(c)	State <b>two</b> structures, visible in Fig. 3.1, that are found only in plant cells.	
	1	
	2	

[4]

[Total: 10]

© UCLES 2017 5090/62/M/J/17