

Cambridge International Examinations Cambridge Ordinary Level

## CHEMISTRY

5070/32 May/June 2017

Paper 3 Practical Test MARK SCHEME Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

® IGCSE is a registered trademark.

## Cambridge O Level – Mark Scheme PUBLISHED

Question		Answer	Marks
1(a)	Titration		12
	<b>Measurements (1)</b> Both readings (initial and final) are present for each titration, readings are recorded to 1dp, no reading is in excess of 50.0 and no initial reading is given as 50.0.		
	<b>Titres (1)</b> All the titres are calculated correctly i.e. no subtraction errors.		
	Accuracy (6) For each of the two best titres give: 3 marks for a titre within 0.2 cm <sup>3</sup> of the Supervisor's value. 2 marks for a titre within 0.3 cm <sup>3</sup> of the Supervisor's value. 1 mark for a titre within 0.4 cm <sup>3</sup> of the Supervisor's value. Concordance (3) Give 3 marks if all the ticked values are within 0.2 cm <sup>3</sup> . Give 2 marks if all the ticked values are within 0.3 cm <sup>3</sup> . Give 1 mark if all the ticked values are within 0.4 cm <sup>3</sup> . Average (1) Circuit 1 mark if the constitute colorulates a correct suprase of colorated titres		
1(b)	Assuming a pipette of 25 cm <sup>3</sup> and the average volume of $\mathbf{Q}$ used = 20.3 cm <sup>3</sup>		
	Moles of sodium thiosulfate	= (20.3 · 0.0230) / 1000 = 0.000467	
1(c)	Moles of iodine	= (b) / 2 = 0.000467 / 2 = 0.000234	1
1(d)	Moles of iodine in 250 cm <sup>3</sup> of <b>P</b>	= (c) · 250 / volume of <b>P</b> used = 0.000234 · 250 / 25 = 0.00234	1

https://xtremepape.rs/

## Cambridge O Level – Mark Scheme PUBLISHED

Question		Answer	Marks		
1(e)	Moles of chlorine in 50 cm <sup>3</sup> of chlorine water	= (d) = 0.00234	1		
1(f)	Mass, in g, of chlorine in 1 dm <sup>3</sup> of chlorine water	= (e) $\cdot$ 71 $\cdot$ 1000 / 50 = 0.00234 $\cdot$ 71 (1) $\cdot$ 1000 / 50 (1) = 3.32	2		
Question 2 General points					
R is ammonium chromium(III) sulfate S is iron(III) chloride					
For gases: to gain credit for the name of the gas produced, the test must be at least partially correct.					
Solutions: colourless is not equivalent to clear and clear is not equivalent to colourless					
No credit is given for conclusions based upon incorrect observations.					
2(a) (test 1)	<ul><li>(a) White ppt (1)</li><li>(b) Insoluble (1)</li></ul>		18		
2(a) (test 2)	Green ppt (1) Insoluble in excess (1)				
2(a) (test 3)	<ul> <li>(a) Green ppt (1) Soluble in excess (1) Green solution (1)</li> <li>(b) Gas turns damp red litmus paper blue (1) Ammonia (1)</li> </ul>				
2(a) (test 4)	(a) White ppt (1) (b) Ppt remains (1)				
2(a) (test 5)	Red-brown ppt (1) Insoluble in excess (1)				

5070/32

## Cambridge O Level – Mark Scheme PUBLISHED

Question	Answer	Marks
2(a) (test 6)	<ul> <li>(a) Yellow colour fades / turns colourless (1)</li> <li>(b) Liquid turns green / black (1) Ppt (1)</li> </ul>	
2(a) (test 7)	<ul> <li>(a) Red–brown solution (1)</li> <li>(b) Liquid turns black-blue (1)</li> </ul>	
2(b)	Conclusions	4
	<b>R</b> contains: ammonium / $NH_4^+$ (1) dependent on a mark being awarded in test 3(b) chromium(III) / $Cr^{3+}$ (1) dependent on insoluble green ppt in test 2 and soluble in test 3 sulfate / $SO_4^{2-}$ (1) dependent on white ppt insoluble in acid in test 1	
	The oxidising agent in <b>S</b> is iron(III) / $Fe^{3+}$ (1)	