

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

	CANDIDATE NAME			
	CENTRE NUMBER		CANDIDATE NUMBER	
* ω	CHEMISTRY			5070/31
00	CHEIMISTRY			5070/31
	Paper 3 Praction	cal Test	October/Novem	ber 2010
4 4 W	-		1 hour 30	minutes
й Л	Candidates ans	swer on the Question Paper		
* 3 8 7 1 4 3 5 2 5 9	Additional Mate	erials: Instructions to Supervisors		
*				

### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black ink.

You may use a soft pencil for any diagrams, graphs or rough work.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Qualitative Analysis Notes are printed on page 8.

You should show the essential steps in any calculations and record experimental results in the spaces provided on the question paper.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
1	
2	
Total	

This document consists of 6 printed pages and 2 blank pages.



1 A large volume of water has been contaminated with acid. Before the contaminated water can be disposed of, the acid must be neutralised by adding calcium carbonate.

For Examiner's Use

 $CaCO_3 + 2H^+ \rightarrow Ca^{2+} + H_2O + CO_2$ 

You are to determine by titration the concentration of hydrogen ions present in the contaminated water and then calculate the mass of calcium carbonate needed to neutralise all the acid.

Solution **P** is a sample of the contaminated water. Solution **Q** is  $0.100 \text{ mol/dm}^3$  sodium hydroxide.

(a) Put P into the burette.

Pipette a  $25.0 \text{ cm}^3$  (or  $20.0 \text{ cm}^3$ ) portion of **Q** into a flask and titrate with **P**, using the indicator provided.

Record your results in the table, repeating the titration as many times as you consider necessary to achieve consistent results.

#### Results

#### Burette readings

titration number	1	2	
final reading / cm <sup>3</sup>			
initial reading / cm <sup>3</sup>			
volume of <b>P</b> used / cm <sup>3</sup>			
best titration results ( $\checkmark$ )			

#### Summary

Tick  $(\checkmark)$  the best titration results.

Using these results, the average volume of **P** required was ...... cm<sup>3</sup>.

Volume of  $\mathbf{Q}$  used was ...... cm<sup>3</sup>.

[12]

© UCLES 2010

https://xtremepape.rs/

(b)	<b>Q</b> is 0.100 mol/dm <sup>3</sup> sodium hydroxide.	For
	Using your results from <b>(a)</b> , calculate the concentration, in mol/dm <sup>3</sup> , of hydrogen ions in <b>P</b> .	Examiner's Use
	concentration of hydrogen ions in <b>P</b> mol/dm <sup>3</sup> [2]	
(c)	The volume of the contaminated water is 10000 dm <sup>3</sup> . Calculate the number of moles of hydrogen ions in this volume of contaminated water.	
	moles of hydrogen ions in 10000 dm <sup>3</sup> of <b>P</b> [1]	
(d)	moles of hydrogen ions in 10000 dm³ of P[1]Using your answer from (c), calculate the minimum mass of calcium carbonate needed to neutralise all the acid in 10000 dm³ of the contaminated water.[1][The relative formula mass of calcium carbonate is 100.]	
	mass of calcium carbonate needed g [2]	
	[Total: 17]	

https://xtremepape.rs/

2 Carry out the following experiments on element **R** and solution **S** and record your observations in the table. You should test and name any gas evolved.

For Examiner's Use

test no.	test	observations
1	To 2 cm depth of dilute hydrochloric acid in a test-tube, add a piece of <b>R</b> and warm the mixture gently until the reaction begins.	
	Once the reaction is complete divide the colourless solution into two equal portions and use them for tests <b>2</b> and <b>3</b> .	
2	To one of the portions of the solution from test <b>1</b> , add aqueous sodium hydroxide until no further change occurs.	
3	To the other portion of the solution from test <b>1</b> , add aqueous ammonia until no further change occurs.	
4	<ul> <li>(a) To 2 cm depth of aqueous sodium hydroxide in a test-tube, add a piece of R and warm the mixture gently until reaction begins. Leave until reaction is complete.</li> </ul>	
	(b) To the colourless solution from (a) add dilute hydrochloric acid until no further change occurs.	

Identify the element R.

**R** is .....

In test 6, R is acting as .....

Solution **S** contains a potassium compound.

Identify the anion in S.

S contains .....

https://xtremepape.rs/

[Total: 23]

[3]

# **BLANK PAGE**

### **BLANK PAGE**

7

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

#### QUALITATIVE ANALYSIS NOTES

# Tests for anions

anion	test	test result
carbonate ( $CO_3^{2-}$ )	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous lead(II) nitrate	yellow ppt.
nitrate (NO $_3^-$ ) [in solution]	add aqueous sodium hydroxide then add aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2–</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous barium nitrate	white ppt.

# Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al <sup>3+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH <sub>4</sub> )	ammonia produced on warming	_
calcium (Ca <sup>2+</sup> )	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper(II) (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

# **Tests for gases**

gas	test and test result
ammonia (NH <sub>3</sub> )	turns damp litmus paper blue
carbon dioxide (CO <sub>2</sub> )	turns limewater milky
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint
oxygen (O <sub>2</sub> )	relights a glowing splint
sulfur dioxide (SO <sub>2</sub> )	turns acidified aqueous potassium dichromate(VI) from orange to green