

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Ordinary Level

	CANDIDATE NAME								
,	CENTRE NUMBER						CANDIDATE NUMBER		
	CHEMISTRY							507	70/31
ມ ມ	Paper 3 Practic	cal Test					Oc	tober/November	2013
								1 hour 30 mii	nutes
	Candidates answer on the Question Paper.								
× μ μ μ μ μ μ μ μ μ μ μ μ μ μ μ μ μ μ μ	Additional Mate	erials:	As liste	d in the C	Confidential	Instructions			
*	READ THESE I	INSTRUC	TIONS I	FIRST					

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black ink.

You may use a soft pencil for any diagrams, graphs or rough work.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

Qualitative Analysis Notes are printed on page 8.

You should show the essential steps in any calculations and record experimental results in the spaces provided on the Question Paper.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
Total		

This document consists of 6 printed pages and 2 blank pages.



1 P is an aqueous solution which contains a mixture of sodium carbonate, Na₂CO₃, and sodium hydroxide, NaOH.

For Examiner's Use

The concentration of sodium carbonate in \mathbf{P} is 0.0200 mol/dm³.

You are to determine by titration the volume of dilute hydrochloric acid, \mathbf{Q} , needed to neutralise a volume of \mathbf{P} , and then calculate the concentration of sodium hydroxide present.

Q is 0.200 mol/dm^3 hydrochloric acid, HCl.

(a) Put Q into the burette.

Pipette a 25.0 cm^3 (or 20.0 cm^3) portion of **P** into a flask and titrate with **Q**, using the indicator provided.

Record your results in the table, repeating the titration as many times as you consider necessary to achieve consistent results.

Results

Burette readings

titration number	1	2	
final reading / cm ³			
initial reading / cm ³			
volume of Q used / cm ³			
best titration results (\checkmark)			

Summary

Tick (\checkmark) the best titration results.

Using these results, the average volume of **Q** required was cm³.

Volume of **P** used was cm³.

[12]

(b) **Q** is $0.200 \text{ mol}/\text{dm}^3$ hydrochloric acid, HC*l*. For Calculate the number of moles of hydrochloric acid present in your average volume Examiner's Use of Q. moles of hydrochloric acid present[1] (c) The concentration of sodium carbonate, Na_2CO_3 , in **P** is 0.0200 mol/dm³. Calculate the number of moles of sodium carbonate present in your volume of P. moles of sodium carbonate present[1] (d) Using your answer to (c), deduce the number of moles of hydrochloric acid which react with the sodium carbonate present in your volume of **P**. $Na_2CO_3 + 2HCl \rightarrow 2NaCl + H_2O + CO_2$ moles of hydrochloric acid which react with the sodium carbonate[1] (e) Using your answers to (b) and (d), calculate the number of moles of hydrochloric acid which react with the sodium hydroxide in your volume of P. moles of hydrochloric acid which react with the sodium hydroxide[1] (f) Using your answer to (e), calculate the concentration, in mol/dm³, of sodium hydroxide in **P**. $NaOH + HCl \rightarrow NaCl + H_2O$ concentration of sodium hydroxide in P mol/dm³ [1] [Total: 17]

2 You are provided with solutions **R** and **S**.

Carry out the following tests and record your observations in the table. You should test and name any gas evolved.

test no.	test	observations
1	(a) To 1 cm depth of R in a test-tube, add an equal volume of aqueous barium nitrate.	
	(b) To the mixture from (a), add dilute nitric acid.	
2	To 2 cm depth of R in a test-tube, add a piece of magnesium ribbon.	
3	To 2 cm depth of R in a test-tube, add a small amount of solid magnesium carbonate.	
4	 (a) To 1 cm depth of S in a test-tube, add a few drops of aqueous silver nitrate. (b) To the mixture from (a), add dilute nitric acid. 	

For Examiner's Use

https://xtremepape.rs/

test no.	test	observations
5 (a)	To 1 cm depth of S in a test-tube, add an equal volume of R . To the mixture, add a little solid manganese(IV) oxide. Mix well then filter off some of the liquid into another test-tube.	
(b)	To the filtrate from (a) , add a few drops of starch solution.	
6 (a)	To 1 cm depth of S in a test-tube, add an equal volume of aqueous copper(II) sulfate.	
(b)	To the mixture from (a) , add an equal volume of R , then a little solid sodium sulfite.	
7 (a)	To 1 cm depth of S in a test-tube, add an equal volume of dilute hydrochloric acid. To the mixture, add a few drops of aqueous potassium iodate(V) solution.	
(b)	To the mixture from (a) , add aqueous sodium hydroxide.	
(c)	To the mixture from (b) , add R .	
		[20]
Conclu	sions	
Identify	a cation and an anion present in R .	
A catior	n present in R is	
An anio	n present in R is	

S contains a sodium compound. Write the formula of the sodium compound in **S**.

The formula of the sodium compound in **S** is[3]

https://xtremepape.rs/

BLANK PAGE

6

BLANK PAGE

7

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

QUALITATIVE ANALYSIS NOTES

Tests for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l⁻</i>) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide then add aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ^{2–}) [in solution]	acidify with dilute nitric acid, then add aqueous barium nitrate	white ppt.

Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al ³⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH ₄ ⁺)	ammonia produced on warming	_
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper(II) (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Tests for gases

gas	test and test result
ammonia (NH ₃)	turns damp litmus paper blue
carbon dioxide (CO ₂)	turns limewater milky
chlorine (Cl ₂)	bleaches damp litmus paper
hydrogen (H ₂)	'pops' with a lighted splint
oxygen (O ₂)	relights a glowing splint