

Cambridge International Examinations

Cambridge Ordinary Level

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

CHEMISTRY 5070/32

Paper 3 Practical Test

October/November 2014

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

Qualitative Analysis Notes are printed on page 8.

You should show the essential steps in any calculations and record experimental results in the spaces provided on the Question Paper.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
Total		

This document consists of 6 printed pages and 2 blank pages.

DC (SJF) 72011/4 © UCLES 2014



[Turn over

1 The active ingredient in bleaching powder is calcium hypochlorite, $Ca(ClO)_2$. When bleaching powder is added to an acidified, aqueous solution of iodide ions, iodine is produced.

$$\mathrm{Ca(C}\mathit{l}\mathrm{O})_{2} \ + \ 4\mathrm{I}^{-} \ + \ 4\mathrm{H}^{+} \ \longrightarrow \ \mathrm{CaC}\mathit{l}_{2} \ + \ 2\mathrm{H}_{2}\mathrm{O} \ + \ 2\mathrm{I}_{2}$$

The amount of iodine produced by the above reaction can be determined by titration with sodium thiosulfate, $Na_2S_2O_3$, using starch as an indicator.

$$2\mathrm{Na}_2\mathrm{S}_2\mathrm{O}_3 \ + \ \mathrm{I}_2 \ \longrightarrow \ \mathrm{Na}_2\mathrm{S}_4\mathrm{O}_6 \ + \ 2\mathrm{NaI}$$

P is an aqueous solution of iodine produced by reacting bleaching powder with an excess of acidified, aqueous iodide ions.

Q is 0.100 mol/dm³ sodium thiosulfate.

(a) Put Q into the burette.

Pipette a 25.0 cm³ (or 20.0 cm³) portion of **P** into a flask.

Add **Q** from the burette until the red-brown colour fades to pale yellow, **then** add a few drops of the starch indicator. This will give a dark blue solution. Continue adding **Q** slowly from the burette until one drop of **Q** causes the blue colour to disappear, leaving a colourless solution. Record your results in the table, repeating the titration as many times as you consider necessary to achieve consistent results.

Results

Burette readings

titration number	1	2	
final reading/cm ³			
initial reading/cm ³			
volume of Q used/cm ³			
best titration results (✓)			

Summary

Tick (\checkmark) the best titration results.

Volume of \mathbf{P} used was cm³. [12]

	3	
(b)	Q is 0.100 mol/dm ³ sodium thiosulfate.	
	Using your results from (a) , calculate the concentration, in mol/dm ³ , of iodine in P .	
	$2Na_2S_2O_3 + I_2 \rightarrow Na_2S_4O_6 + 2NaI$	
	concentration of iodine in P mol/dm ³ [2]	
(c)	Using your answer from (b) , deduce the number of moles of calcium hypochlorite required to produce the iodine in 1 dm ³ of P .	
	$\mathrm{Ca(C}l\mathrm{O)_2} \ + \ 4\mathrm{I^-} \ + \ 4\mathrm{H^+} \ \longrightarrow \ \mathrm{CaC}l_2 \ + \ 2\mathrm{H_2O} \ + \ 2\mathrm{I_2}$	
	moles of calcium hypochlorite[1]	
(d)	Given that the number of moles of calcium hypochlorite in your answer from (c) were present in 10.0 g of the bleaching powder, calculate the percentage by mass of calcium hypochlorite in the bleaching powder. [The relative formula mass of $Ca(ClO)_2$ is 143.]	

percentage by mass of calcium hypochlorite in the bleaching powder% [2]

You are provided with solutions **R** and **S**.

Carry out the following tests and record your observations in the table. You should test and name any gas evolved.

test no.	test	observations
1	Gently warm 2 cm depth of R in a test-tube.	
2	 (a) To 1 cm depth of aqueous sodium chloride in a test-tube, add a few drops of aqueous silver nitrate. (b) To the mixture from (a), add R until no further change occurs. 	
3	To 1 cm depth of aqueous copper(II) sulfate in a test-tube, add R until no further change occurs.	
4	To 1 cm depth of S in a test-tube, add R until no further change occurs. Keep this mixture for use in Test 5.	
5	To 1 cm depth of aqueous hydrogen peroxide in a boiling tube, add the mixture from Test 4.	

test no.		test	observations
6	(a)	To 1 cm depth of S in a test-tube, add an equal volume of aqueous silver nitrate. Leave to settle.	
	(b)	To the mixture from (a) , add dilute nitric acid.	
7	(a)	To 1 cm depth of S in a test-tube, add an equal volume of aqueous sodium thiosulfate and mix well.	
	(b)	To the mixture from (a) , add R until no further change occurs.	

[19]

Conclusions

[Total: 23]

BLANK PAGE

BLANK PAGE

QUALITATIVE ANALYSIS NOTES

Tests for anions

anion	test	test result
carbonate (CO ₃ ²⁻)	add dilute acid	effervescence, carbon dioxide produced
chloride (C <i>l</i> ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO ₃ ⁻) [in solution]	add aqueous sodium hydroxide then add aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify with dilute nitric acid, then add aqueous barium nitrate	white ppt.

Tests for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia	
aluminium (Al ³⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess	
ammonium (NH ₄ ⁺)	ammonia produced on warming	_	
calcium (Ca ²⁺)	white ppt., insoluble in excess	no ppt., or very slight white ppt.	
copper(II) (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution	
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess	
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess	
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution	

Tests for gases

gas	test and test result		
ammonia (NH ₃)	turns damp red litmus paper blue		
carbon dioxide (CO ₂)	turns limewater milky		
chlorine (Cl ₂)	bleaches damp litmus paper		
hydrogen (H ₂)	'pops' with a lighted splint		
oxygen (O ₂)	relights a glowing splint		

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.