MARK SCHEME for the May/June 2012 question paper

for the guidance of teachers

4037 ADDITIONAL MATHEMATICS

4037/22

Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2012 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE O LEVEL – May/June 2012	4037	22

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Accuracy mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
 B2, 1, 0 means that the candidate can earn anything from 0 to 2.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE O LEVEL – May/June 2012	4037	22

The following abbreviations may be used in a mark scheme or used on the scripts:

- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)

Penalties

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures this is regarded as an error in accuracy.
- OW –1, 2 This is deducted from A or B marks when essential working is omitted.
- PA –1 This is deducted from A or B marks in the case of premature approximation.
- S –1 Occasionally used for persistent slackness usually discussed at a meeting.
- EX –1 Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

	Page 4		: Teachers' version	Syllabus	Paper
		GCE O LEVE	L – May/June 2012	4037	22
1	(i) 7 ∈ <i>P</i>				B1
	(ii) 8 ∉ <i>S</i>				B1
	(iii) $n(N \cap S)$) = 6			B1
2	(i) $k\sqrt{4x+1}$ k=6 at	oe llow unsimplified			M1 A1
	(ii) Use $\partial y =$	$\frac{dy}{dt} \times p$			M1
	30 <i>p</i>	$ax_{(x=6)}$			A1√
3		or $9b^2 - 4ac \approx 0$	OR Eliminate y and m $((2x+3)x-5 = x^2 + 3x + 4)$ Solve quadratic for x		M1 A1 M1
	Solve for 2 val		Solve for 2 values of <i>m</i>		M1
	Solve for 2 val	lues of <i>m</i>			A1
4	(i) $\begin{pmatrix} 4 & 1 & 7 \\ 2 & 5 & 1 \end{pmatrix}$	$ \begin{pmatrix} 5\\3\\1 \end{pmatrix} \text{or transpose} $ $ \begin{pmatrix} 2\\6 \end{pmatrix} \begin{pmatrix} 8\\4\\2 \end{pmatrix} \text{or transpose} $			B1
	$+\begin{pmatrix}2&5\\4&3\end{pmatrix}$	$ \binom{2}{6} \binom{8}{4} $ or transpose			B1+B1
	(ii) $\begin{pmatrix} 30\\ 26 \end{pmatrix}$ or	$\begin{pmatrix} 40\\56 \end{pmatrix}$ or $\begin{pmatrix} 30\\y \end{pmatrix}$ and $\begin{pmatrix} 40\\y \end{pmatrix}$	$\begin{pmatrix} 40 \\ y \end{pmatrix}$ or $\begin{pmatrix} x \\ 26 \end{pmatrix}$ and $\begin{pmatrix} x \\ 56 \end{pmatrix}$ from	n correct part (i)	B1
	Claire 70	and Denise 82			B1
5	(i) $f(2)(=8)$ k = 4	+4k-16-8)=0			M1 A1
	$x^2 + 6x +$ Use quade	ratic formula or completir	ng square		M1 A1 M1
	$\frac{-6\pm\sqrt{6^2}}{}$	$\frac{-4 \times (1) \times 4}{2}$			A1
	$-3\pm\sqrt{5}$	2			B1√

	Page 5		Scheme: Teachers' version	Syllabus	Paper
		GCE	O LEVEL – May/June 2012	4037	22
6	(a) (i) (-2)	³ or 35			B1
	-280				B1
	(ii) $2^2 \times 2^2$	$21(x^2)$			B1
		$-280) + 4 \times (84)$			M1
	-504	, , ,			M1 A1
	-304				111
		$(3)^{2}$			
	(b) Identify x	$z^4 \times \left(\frac{z}{x^2}\right)$			B1
	×15 only				B1
	•				
	$135(x^0)$				B1
7	(i) $\ln y = \ln y$	$a + b \ln x$ OR	$\lg y = \lg a + b \lg x$ may be implied		B1
/	• •				M1
		61 3.40	$x / \lg x$ with attempt at linear scale 5.01 5.99		
		.19 3.09	3.89 4.39		A2, 1, 0
	$\lim y = 2.$	5.07	J.07 T.J7		
	$\lg x = 0.$	70 1.48	2.18 2.60		
	•	95 1.34	1.69 1.91		
	-87 0.				
	(Marks fo	r points and line	on graph NOT for table)		
		_			
	(ii) Calculates $b = 0.5 \pm$		ight line log graph		M1 A1
			straight line log graph		M1
	$a = 4 \pm 0$	•	straight line log graph		A1
		able graph or for	mula		M1
	32 to 49				A1
8	$x^2h = 256$		OR		B1
0	$x n = 230$ $A = x^2 + 4xh$				M1
			$A = \frac{256}{h} + 64\sqrt{h}$		
	$A = x^2 + \frac{1024}{x}$				A1
	dA = 102	4	$\frac{\mathrm{d}A}{\mathrm{d}h} = \frac{-256}{h^2} + \frac{32}{\sqrt{h}}$		A1√
	$\frac{\mathrm{d}A}{\mathrm{d}x} = 2x - \frac{102}{x^2}$	– oe	$dh h^2 \sqrt{h}$		
	Equate to 0 and		h = 4		M1
	x = 8		<i>x</i> = 8		A1
	h = 4				A1

Page 6		ige 6		Paper
			GCE O LEVEL – May/June 2012 4037	22
9	(a)	(i)	$\tan x = \frac{5}{3}$	B1
			x = 59(.0)	B1
			x = 239(.0) and no others	B1
		(ii)	Use $\sin^2 y = 1 - \cos^2 y$	B1
			$5\cos^2 y - 9\cos y - 2 = 0$	B1
			Solve 3 term quadratic (in cosy)	M1
			101.5	A1
			258.5 and no others	B1
	(b)	(3-	z) = 0.927 or 0.93	B1
		2.07		B1
		(3-	$z) = \pi - 0.927$	M1
		0.78	6 or 0.785 or 0.79 and no others	A1
10	(a)	(i)	792	B1
		(ii)	4W, 3M and 5W, 2M	M1
			5×35 or (1) $\times 21$	B1
			196	A1
	(b)	(i)	$4 \times 5 \times 4 \times 3$ or $\frac{2}{3} \times 6 \times 5 \times 4 \times 3$	M1
			240	A1
			1 (2.2)	
		(ii)	$4 \times 4 \times 3 \times 1$ or $\frac{1}{5} \times (240)$	M1
			48	A1

Page 7		7	Mark Scheme: Teachers' version	Syllabus	Paper
			GCE O LEVEL – May/June 2012	4037	22
11 E	(i)		$k\cos\frac{1}{2}x \left(\frac{1}{2}\cos\frac{1}{2}x\right)$		M1
			ient tangent $-\frac{1}{4}\sqrt{2}$ or -0.35		A1
			$\frac{\sqrt{2}}{2} = -\frac{1}{4}\sqrt{2}\left(x - \frac{3\pi}{2}\right)$		M1
		<i>y</i> = 0	$x = \frac{3\pi}{2} + 2 \text{ or } 6.71$		A1
	(ii)	MET	HOD A		
		∫sin	$\frac{1}{2}xdx = -2\cos\frac{1}{2}x$		B1
			ify 2π	X	B1
			imits of 1.5 π and (2π) on $k \cos \frac{1}{2}x$ $\left(2 - \sqrt{2} \text{ or } 0.58\right)$	36)	M1
		Atter	npt at area of triangle $\left(=\frac{\sqrt{2}}{2}=0.707\right)$		M1
		Plan	of area of triangle subtract area under curve. completely correct		M1 M1
		$\frac{3\sqrt{2}}{2}$	-2 or 0.121		A1
		MET	HOD B		
			g integral of (equation of line $-$ equation of curve)		M1
		Jsin	$\frac{1}{2}xdx = -2\cos\frac{1}{2}x$		B1
			ify 2π		B1
			imits of 1.5π and (2π) on $k\cos\frac{1}{2}x$		M1
		Use 1	imits of 1.5π and (x_Q) on integral of equation of line		M1
			completely correct		M1
		$\frac{3\sqrt{2}}{2}$	-2 or 0.121		A1

Page 8	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE O LEVEL – May/June 2012	4037	22
l O (i) Uses	product rule		M1
(1	$(x)e^{-x}$		A1
∫(1-	$-x)e^{-x}dx = xe^{-x}$		M1
∫xe⁻	$\int dx = -xe^{-x} + \int e^{-x} dx = -xe^{-x} - e^{-x}$		Alag
(ii) gradi	ent tangent $= -\frac{1}{e^2}$ or $= -0.135$		B1
<i>y</i> – -	$\frac{2}{e^2} = -\frac{1}{e^2}(x-2)$		M1
Uses	line cuts y-axis at $\frac{4}{e^2}$ or 0.541		A1
Area	trapezium $\left(= \frac{6}{e^2} \text{ or } 0.812 \right)$		M1
Uses	limits of 2 and 0 on $-xe^{-x} - e^{-x}$ (=1 $-\frac{3}{e^2}$ or 0.594)		M1
_	ate area of trapezium subtract area under curve		M1
$\frac{9}{e^2}$ –	1 or 0.218		A1