

Mathematical Formulae**1. ALGEBRA***Quadratic Equation*

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} .$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \dots + \binom{n}{r} a^{n-r} b^r + \dots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.

2. TRIGONOMETRY*Identities*

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$


Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} .$$

$$a^2 = b^2 + c^2 - 2bc \cos A.$$

$$\Delta = \frac{1}{2} bc \sin A.$$

1



- (i) Copy the Venn diagram above and shade the region that represents $A \cup (B \cap C)$. [1]
- (ii) Copy the Venn diagram above and shade the region that represents $A \cap (B \cup C)$. [1]
- (iii) Copy the Venn diagram above and shade the region that represents $(A \cup B \cup C)'$. [1]
- 2 Find the set of values of x for which $(2x + 1)^2 > 8x + 9$. [4]
- 3 Prove that $\frac{\sin A}{1 + \cos A} + \frac{1 + \cos A}{\sin A} \equiv 2 \operatorname{cosec} A$. [4]
- 4 A function f is such that $f(x) = ax^3 + bx^2 + 3x + 4$. When $f(x)$ is divided by $x - 1$, the remainder is 3. When $f(x)$ is divided by $2x + 1$, the remainder is 6. Find the value of a and of b . [5]
- 5 Given that $\mathbf{a} = 5\mathbf{i} - 12\mathbf{j}$ and that $\mathbf{b} = p\mathbf{i} + \mathbf{j}$, find
- (i) the unit vector in the direction of \mathbf{a} , [2]
- (ii) the values of the constants p and q such that $q\mathbf{a} + \mathbf{b} = 19\mathbf{i} - 23\mathbf{j}$. [3]
- 6 (i) Solve the equation $2t = 9 + \frac{5}{t}$. [3]
- (ii) Hence, or otherwise, solve the equation $2x^{\frac{1}{2}} = 9 + 5x^{-\frac{1}{2}}$. [3]
- 7 (i) Express $4x^2 - 12x + 3$ in the form $(ax + b)^2 + c$, where a , b and c are constants and $a > 0$. [3]
- (ii) Hence, or otherwise, find the coordinates of the stationary point of the curve $y = 4x^2 - 12x + 3$. [2]
- (iii) Given that $f(x) = 4x^2 - 12x + 3$, write down the range of f . [1]

- 8 A curve is such that $\frac{d^2y}{dx^2} = 4e^{-2x}$. Given that $\frac{dy}{dx} = 3$ when $x = 0$ and that the curve passes through the point $(2, e^{-4})$, find the equation of the curve. [6]

- 9 (i) Find, in ascending powers of x , the first 3 terms in the expansion of $(2 - 3x)^5$. [3]

The first 3 terms in the expansion of $(a + bx)(2 - 3x)^5$ in ascending powers of x are $64 - 192x + cx^2$.

- (ii) Find the value of a , of b and of c . [5]

- 10 (a) Functions f and g are defined, for $x \in \mathbb{R}$, by

$$f(x) = 3 - x,$$

$$g(x) = \frac{x}{x+2}, \text{ where } x \neq -2.$$

- (i) Find $fg(x)$. [2]

- (ii) Hence find the value of x for which $fg(x) = 10$. [2]

- (b) A function h is defined, for $x \in \mathbb{R}$, by $h(x) = 4 + \ln x$, where $x > 1$.

- (i) Find the range of h . [1]

- (ii) Find the value of $h^{-1}(9)$. [2]

- (iii) On the same axes, sketch the graphs of $y = h(x)$ and $y = h^{-1}(x)$. [3]

- 11 Solve the equation

- (i) $\tan 2x - 3 \cot 2x = 0$, for $0^\circ < x < 180^\circ$, [4]

- (ii) $\operatorname{cosec} y = 1 - 2 \cot^2 y$, for $0^\circ \leq y \leq 360^\circ$, [5]

- (iii) $\sec\left(z + \frac{\pi}{2}\right) = -2$, for $0 < z < \pi$ radians. [3]

12 Answer only **one** of the following two alternatives.

EITHER

A curve has equation $y = \frac{x^2}{x+1}$.

- (i) Find the coordinates of the stationary points of the curve. [5]

The normal to the curve at the point where $x = 1$ meets the x -axis at M . The tangent to the curve at the point where $x = -2$ meets the y -axis at N .

- (ii) Find the area of the triangle MNO , where O is the origin. [6]

OR

A curve has equation $y = e^{x-2} - 2x + 6$.

- (i) Find the coordinates of the stationary point of the curve and determine the nature of the stationary point. [6]

The area of the region enclosed by the curve, the positive x -axis, the positive y -axis and the line $x = 3$ is $k + e - e^{-2}$.

- (ii) Find the value of k . [5]

BLANK PAGE

BLANK PAGE

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.